

Diagnoses of HIV and/or AIDS Reported in Stark County

Diagnoses of HIV infection as of June 30, 2014, by disease status at initial diagnosis and by selected characteristics, Stark County

Characteristic	2013 diagnosis of HIV infection			Disease Status at Initial Diagnosis					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	7.2	13	76%	11	73%	1	100%	1	100%
Females	*	4	24%	4	27%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	*	4	24%	4	27%	-	-	-	-
25-29	*	3	18%	3	20%	-	-	-	-
30-34	*	3	18%	2	13%	-	-	1	100%
35-39	*	-	-	-	-	-	-	-	-
40-44	*	1	6%	1	7%	-	-	-	-
45-49	*	3	18%	3	20%	-	-	-	-
50-54	*	1	6%	1	7%	-	-	-	-
55-64	*	2	12%	1	7%	1	100%	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity									
White, not Hispanic	4.0	13	76%	11	73%	1	100%	1	100%
Black/African American, not Hispanic	*	4	24%	4	27%	-	-	-	-
Hispanic/Latino	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Race/Sex									
White, not Hispanic Males	5.7	9	53%	7	47%	1	100%	1	100%
White, not Hispanic Females	*	4	24%	4	27%	-	-	-	-
Black/African American, not Hispanic Males	*	4	24%	4	27%	-	-	-	-
Black/African American, not Hispanic Females	*	-	-	-	-	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-	-	-
Total	4.5	17		15		1		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Diagnoses of HIV infection as of June 30, 2014, by disease status at initial diagnosis and by transmission category, Stark County

Transmission Category ^a	2013 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	9	69%	8	73%	-	-	1	100%
Injection drug use (IDU)	1	8%	1	9%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	3	23%	2	18%	1	100%	-	-
Subtotal	13	100%	11	100%	1	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	1	25%	1	25%	-	-	-	-
Other/unknown	3	75%	3	75%	-	-	-	-
Subtotal	4	100%	4	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	17		15		1		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Diagnoses of HIV infection as of June 30, 2014, by disease status at initial diagnosis and by exposure category, Stark County

Exposure Category ^a	2013 diagnosis of HIV infection		Disease Status at Initial Diagnosis					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	9	53%	8	53%	-	-	1	100%
Injection drug use (IDU) only	1	6%	1	7%	-	-	-	-
Heterosexual contact only	1	6%	1	7%	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	6	35%	5	33%	1	100%	-	-
Total	17		15		1		1	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS).

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Diagnoses of HIV infection as of June 30, 2014, by race/ethnicity and transmission category, Stark County

Transmission Category ^a	2013 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	6	67%	3	75%	-	-	-	-	-	-	-	-
Injection drug use (IDU)	1	11%	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	2	22%	1	25%	-	-	-	-	-	-	-	-
Subtotal	9	100%	4	100%	-	-	-	-	-	-	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	25%	-	-	-	-	-	-	-	-	-	-
Other/unknown	3	75%	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	-	-	-	-	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	13		4		0		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Diagnoses of HIV infection as of June 30, 2014, by race/ethnicity and exposure category, Stark County

Exposure Category ^a	2013 Diagnosis of HIV Infection											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	6	46%	3	75%	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	1	8%	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	1	8%	-	-	-	-	-	-	-	-	-	-
MSM & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	5	38%	1	25%	-	-	-	-	-	-	-	-
Total	13		4		0		0		0		0	

Note: Living with a diagnosis of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Diagnoses of HIV infection as of June 30, 2014, by year of diagnosis and by selected characteristics, Stark County

Diagnosis of HIV Infection by Year

Characteristic	2009			2010			2011			2012		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex												
Males	12.1	22	85%	13.8	25	89%	13.8	25	86%	7.2	13	87%
Females	*	4	15%	*	3	11%	*	4	14%	*	2	13%
Age at diagnosis (yr)												
<13	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	4%	*	4	14%	*	2	7%	*	1	7%
20-24	37.7	9	35%	*	3	11%	33.5	8	28%	*	-	-
25-29	*	1	4%	*	2	7%	23.9	5	17%	*	3	20%
30-34	*	3	12%	*	3	11%	*	2	7%	*	2	13%
35-39	*	3	12%	24.8	5	18%	*	2	7%	*	-	-
40-44	*	2	8%	*	4	14%	*	1	3%	*	4	27%
45-49	*	3	12%	*	1	4%	*	2	7%	*	3	20%
50-54	*	2	8%	*	4	14%	*	2	7%	*	2	13%
55-64	*	2	8%	*	2	7%	*	4	14%	*	-	-
65+	*	-	-	*	-	-	*	1	3%	*	-	-
Race/Ethnicity												
White, not Hispanic	5.5	18	69%	5.5	18	64%	5.5	18	62%	3.7	12	80%
Black/African American, not Hispanic	17.6	5	19%	28.1	8	29%	31.6	9	31%	*	3	20%
Hispanic/Latino	*	-	-	*	1	4%	*	-	-	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	12%	*	1	4%	*	2	7%	*	-	-
Race/Sex												
White, not Hispanic Males	10.7	17	65%	10.1	16	57%	10.1	16	55%	6.3	10	67%
White, not Hispanic Females	*	1	4%	*	2	7%	*	2	7%	*	2	13%
Black/African American, not Hispanic Males	*	3	12%	50.1	7	25%	50.1	7	24%	*	3	20%
Black/African American, not Hispanic Females	*	2	8%	*	1	4%	*	2	7%	*	-	-
Hispanic/Latino Males	*	-	-	*	1	4%	*	-	-	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	3	12%	*	1	4%	*	2	7%	*	-	-
Total	6.9	26		7.5	28		7.7	29		4.0	15	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2009-2012) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Diagnoses of HIV infection as of June 30, 2014, by year of diagnosis and by transmission category, Stark County

Diagnosis of HIV Infection by Year

Transmission Category ^a	2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	17	77%	19	76%	20	80%	11	85%
Injection drug use (IDU)	1	5%	-	-	1	4%	-	-
Male-to-male sexual contact and IDU	1	5%	2	8%	-	-	1	8%
Heterosexual contact	3	14%	3	12%	2	8%	-	-
Other/unknown	-	-	1	4%	2	8%	1	8%
Subtotal	22	100%	25	100%	25	100%	13	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	1	25%	-	-
Heterosexual contact	4	100%	3	100%	3	75%	1	50%
Other/unknown	-	-	-	-	-	-	1	50%
Subtotal	4	100%	3	100%	4	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	26		28		29		15	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2009-2012) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Diagnoses of HIV infection as of June 30, 2014, by year of diagnosis and by exposure category, Stark County

Diagnosis of HIV Infection by Year

Exposure Category ^a	2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	16	62%	18	64%	18	62%	11	73%
Injection drug use (IDU) only	1	4%	-	-	1	3%	-	-
Heterosexual contact only	7	27%	6	21%	5	17%	1	7%
MSM & IDU	1	4%	1	4%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	3%	-	-
MSM & Heterosexual contact	1	4%	1	4%	2	7%	-	-
MSM & IDU & Heterosexual contact	-	-	1	4%	-	-	1	7%
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	4%	2	7%	2	13%
Total	26		28		29		15	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2009-2012) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported HIV (not AIDS) diagnoses as of June 30, 2013, by year of HIV diagnosis and by selected characteristics, Stark County

Characteristic	Year of HIV (not AIDS) Diagnosis									
	2009		2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	14	82%	18	90%	18	90%	12	86%	11	73%
Females	3	18%	2	10%	2	10%	2	14%	4	27%
Age at HIV diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	1	6%	4	20%	2	10%	1	7%	-	-
20-24	9	53%	2	10%	8	40%	-	-	4	27%
25-29	1	6%	1	5%	3	15%	3	21%	3	20%
30-34	2	12%	3	15%	2	10%	2	14%	2	13%
35-39	1	6%	3	15%	1	5%	-	-	-	-
40-44	-	-	1	5%	1	5%	4	29%	1	7%
45-49	2	12%	1	5%	-	-	2	14%	3	20%
50-54	-	-	4	20%	1	5%	2	14%	1	7%
55-64	1	6%	1	5%	2	10%	-	-	1	7%
65+	-	-	-	-	-	-	-	-	-	-
Race/Ethnicity										
White, not Hispanic	12	71%	14	70%	16	80%	11	79%	11	73%
Black/African American, not Hispanic	3	18%	4	20%	3	15%	3	21%	4	27%
Hispanic/Latino	-	-	1	5%	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	2	12%	1	5%	1	5%	-	-	-	-
Race/Sex										
White, not Hispanic Males	11	65%	12	60%	14	70%	9	64%	7	47%
White, not Hispanic Females	1	6%	2	10%	2	10%	2	14%	4	27%
Black/African American, not Hispanic Males	2	12%	4	20%	3	15%	3	21%	4	27%
Black/African American, not Hispanic Females	1	6%	-	-	-	-	-	-	-	-
Hispanic/Latino Males	-	-	1	5%	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	2	12%	1	5%	1	5%	-	-	-	-
Total	17		20		20		14		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2012.

Reported HIV (not AIDS) diagnoses as of June 30, 2014, by year of HIV diagnosis and by transmission category, Stark County

Transmission Category ^a	Year of HIV (not AIDS) Diagnosis									
	2009		2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	13	93%	13	72%	15	83%	11	92%	8	73%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	9%
Male-to-male sexual contact and IDU	-	-	2	11%	-	-	-	-	-	-
Heterosexual contact	1	7%	2	11%	1	6%	-	-	-	-
Other/unknown	-	-	1	6%	2	11%	1	8%	2	18%
Subtotal	14	100%	18	100%	18	100%	12	100%	11	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	3	100%	2	100%	2	100%	1	50%	1	25%
Other/unknown	-	-	-	-	-	-	1	50%	3	75%
Subtotal	3	100%	2	100%	2	100%	2	100%	4	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	17		20		20		14		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported HIV (not AIDS) diagnoses as of June 30, 2014, by year of HIV diagnosis and by exposure category, Stark County

Exposure Category ^a	Year of HIV (not AIDS) Diagnosis									
	2009		2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	12	71%	12	60%	14	70%	11	79%	8	53%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	1	7%
Heterosexual contact only	4	24%	4	20%	3	15%	1	7%	1	7%
MSM & IDU	-	-	1	5%	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	1	6%	1	5%	1	5%	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	1	5%	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	5%	2	10%	2	14%	5	33%
Total	17		20		20		14		15	

Note: Reported HIV (not AIDS) diagnoses include persons diagnosed with HIV (not AIDS) in the given calendar year. Cases could have progressed to AIDS in a subsequent calendar year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported cases of AIDS as of June 30, 2014, by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Stark County

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2009		2010		2011		2012		2013		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	11	85%	7	64%	11	79%	3	100%	4	100%	302	84%
Females	2	15%	4	36%	3	21%	-	-	-	-	56	16%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	2%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	5	1%
20-24	2	15%	1	9%	1	7%	-	-	1	25%	37	10%
25-29	-	-	2	18%	3	21%	-	-	-	-	53	15%
30-34	3	23%	2	18%	1	7%	-	-	2	50%	71	20%
35-39	1	8%	1	9%	1	7%	2	67%	-	-	64	18%
40-44	2	15%	4	36%	1	7%	-	-	-	-	48	13%
45-49	2	15%	-	-	3	21%	1	33%	-	-	32	9%
50-54	2	15%	-	-	1	7%	-	-	-	-	20	6%
55-64	1	8%	1	9%	2	14%	-	-	1	25%	17	5%
65+	-	-	-	-	1	7%	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	10	77%	4	36%	4	29%	2	67%	2	50%	251	70%
Black/African American, not Hispanic	2	15%	6	55%	9	64%	1	33%	2	50%	97	27%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	6	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	8%	1	9%	1	7%	-	-	-	-	4	1%
Race/Sex												
White, not Hispanic Males	8	62%	4	36%	4	29%	2	67%	2	50%	222	62%
White, not Hispanic Females	2	15%	-	-	-	-	-	-	-	-	29	8%
Black/African American, not Hispanic Males	2	15%	2	18%	6	43%	1	33%	2	50%	70	20%
Black/African American, not Hispanic Females	-	-	4	36%	3	21%	-	-	-	-	27	8%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	6	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	8%	1	9%	1	7%	-	-	-	-	4	1%
Total	13		11		14		3		4		358	

Note: AIDS diagnoses by year (2009-2013) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2013.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported cases of AIDS as of June 30, 2014, by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Stark County

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2009		2010		2011		2012		2013		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	5	45%	6	86%	7	64%	2	67%	3	75%	215	72%
Injection drug use (IDU)	1	9%	-	-	1	9%	-	-	-	-	13	4%
Male-to-male sexual contact and IDU	1	9%	-	-	-	-	1	33%	-	-	13	4%
Heterosexual contact	2	18%	-	-	2	18%	-	-	-	-	16	5%
Other/unknown	2	18%	1	14%	1	9%	-	-	1	25%	43	14%
Subtotal	11	100%	7	100%	11	100%	3	100%	4	100%	300	100%
Female adult or adolescent												
Injection drug use	1	50%	-	-	1	33%	-	-	-	-	13	25%
Heterosexual contact	1	50%	4	100%	2	67%	-	-	-	-	34	65%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	10%
Subtotal	2	100%	4	100%	3	100%	-	-	-	-	52	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	13		11		14		3		4		358	

Note: AIDS diagnoses by year (2009-2013) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2013.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported cases of AIDS as of June 30, 2014, by year of AIDS diagnosis and cumulative AIDS diagnoses by exposure category, Stark County

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2009		2010		2011		2012		2013		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	5	38%	6	55%	6	43%	2	67%	3	75%	208	58%
Injection drug use (IDU) only	1	8%	-	-	1	7%	-	-	-	-	15	4%
Heterosexual contact only	3	23%	4	36%	4	29%	-	-	-	-	50	14%
MSM & IDU	1	8%	-	-	-	-	-	-	-	-	7	2%
IDU & Heterosexual contact	1	8%	-	-	1	7%	-	-	-	-	11	3%
MSM & Heterosexual contact	-	-	-	-	1	7%	-	-	-	-	7	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	1	33%	-	-	6	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	6	2%
Other/unknown	2	15%	1	9%	1	7%	-	-	1	25%	48	13%
Total	13		11		14		3		4		358	

Note: AIDS diagnoses by year (2009-2013) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2013.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Persons Living with a Diagnosis of HIV and/or AIDS Reported in Stark County

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by current disease status and by selected characteristics, Stark County

Characteristic	Living with a diagnosis of HIV infection in 2013			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	161.4	293	81%	174	81%	119	82%
Females	35.2	68	19%	42	19%	26	18%
Age at end of year							
<13	*	4	1%	2	1%	2	1%
13-14	*	-	-	-	-	-	-
15-19	75.7	19	5%	16	7%	3	2%
20-24	247.4	59	16%	45	21%	14	10%
25-29	253.6	53	15%	31	14%	22	15%
30-34	290.2	62	17%	34	16%	28	19%
35-39	283.1	57	16%	32	15%	25	17%
40-44	169.6	41	11%	19	9%	22	15%
45-49	109.0	28	8%	16	7%	12	8%
50-54	69.2	20	6%	12	6%	8	6%
55-64	30.1	16	4%	9	4%	7	5%
65+	*	2	1%	-	-	2	1%
Race/Ethnicity							
White, not Hispanic	76.5	251	70%	157	73%	94	65%
Black/African American, not Hispanic	334.0	95	26%	50	23%	45	31%
Hispanic/Latino	107.6	7	2%	4	2%	3	2%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	8	2%	5	2%	3	2%
Race/Sex							
White, not Hispanic Males	133.1	211	58%	128	59%	83	57%
White, not Hispanic Females	23.6	40	11%	29	13%	11	8%
Black/African American, not Hispanic Males	487.1	68	19%	38	18%	30	21%
Black/African American, not Hispanic Females	186.5	27	7%	12	6%	15	10%
Hispanic/Latino Males	204.5	7	2%	4	2%	3	2%
Hispanic/Latina Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	8	2%	5	2%	3	2%
Total	96.3	361		216		145	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2013.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2012 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by current disease status and by transmission category, Stark County

Transmission Category ^a	Living with a diagnosis of HIV infection in 2013		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	213	73%	127	73%	86	72%
Injection drug use (IDU)	3	1%	2	1%	1	1%
Male-to-male sexual contact and IDU	9	3%	4	2%	5	4%
Heterosexual contact	16	5%	7	4%	9	8%
Other/unknown	52	18%	34	20%	18	15%
Subtotal	293	100%	174	100%	119	100%
Female adult or adolescent						
Injection drug use	6	9%	3	8%	3	13%
Heterosexual contact	41	64%	22	55%	19	79%
Other/unknown	17	27%	15	38%	2	8%
Subtotal	64	100%	40	100%	24	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	100%	2	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	4	100%	2	100%	2	100%
Total	361		216		145	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2013.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by current disease status and by exposure category, Stark County

Exposure Category ^a	Living with a diagnosis of HIV infection in 2013		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	204	57%	122	56%	82	57%
Injection drug use (IDU) only	6	2%	3	1%	3	2%
Heterosexual contact only	57	16%	29	13%	28	19%
MSM & IDU	4	1%	3	1%	1	1%
IDU & Heterosexual contact	3	1%	2	1%	1	1%
MSM & Heterosexual contact	9	2%	5	2%	4	3%
MSM & IDU & Heterosexual contact	5	1%	1	<1%	4	3%
Perinatal exposure	4	1%	2	1%	2	1%
Other/unknown	69	19%	49	23%	20	14%
Total	361		216		145	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2013.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by race/ethnicity and by transmission category, Stark County

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2013											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	162	77%	45	66%	4	57%	-	-	-	-	2	29%
Injection drug use (IDU)	2	1%	1	1%	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	7	3%	2	3%	-	-	-	-	-	-	-	-
Heterosexual contact	5	2%	6	9%	1	14%	-	-	-	-	4	57%
Other/unknown	35	17%	14	21%	2	29%	-	-	-	-	1	14%
Subtotal	211	100%	68	100%	7	100%	-	-	-	-	7	100%
Female adult or adolescent												
Injection drug use	4	11%	2	8%	-	-	-	-	-	-	-	-
Heterosexual contact	21	55%	19	76%	-	-	-	-	-	-	1	100%
Other/unknown	13	34%	4	16%	-	-	-	-	-	-	-	-
Subtotal	38	100%	25	100%	-	-	-	-	-	-	1	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	2	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	2	100%	-	-	-	-	-	-	-	-
Total	251		95		7		0		0		8	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2013.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by race/ethnicity and by exposure category, Stark County

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2013											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	158	63%	41	43%	4	57%	-	-	-	-	1	13%
Injection drug use (IDU) only	4	2%	2	2%	-	-	-	-	-	-	-	-
Heterosexual contact only	26	10%	25	26%	1	14%	-	-	-	-	5	63%
MSM & IDU	4	2%	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	2	1%	1	1%	-	-	-	-	-	-	-	-
MSM & Heterosexual contact	4	2%	4	4%	-	-	-	-	-	-	1	13%
MSM & IDU & Heterosexual contact	3	1%	2	2%	-	-	-	-	-	-	-	-
Perinatal exposure	2	1%	2	2%	-	-	-	-	-	-	-	-
Other/unknown	48	19%	18	19%	2	29%	-	-	-	-	1	13%
Total	251		95		7		0		0		8	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2013.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by year and by selected characteristics, Stark County

Living with a Diagnosis of HIV Infection

Characteristic	2009		2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	221	80%	246	81%	268	81%	281	81%	293	81%
Females	55	20%	58	19%	62	19%	64	19%	68	19%
Age at end of year										
<13	4	1%	4	1%	4	1%	4	1%	4	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	12	4%	16	5%	18	5%	19	6%	19	5%
20-24	45	16%	48	16%	56	17%	56	16%	59	16%
25-29	40	14%	42	14%	47	14%	50	14%	53	15%
30-34	52	19%	55	18%	57	17%	59	17%	62	17%
35-39	50	18%	55	18%	57	17%	57	17%	57	16%
40-44	31	11%	35	12%	36	11%	40	12%	41	11%
45-49	19	7%	20	7%	22	7%	25	7%	28	8%
50-54	12	4%	16	5%	17	5%	19	6%	20	6%
55-64	10	4%	12	4%	14	4%	14	4%	16	4%
65+	1	<1%	1	<1%	2	1%	2	1%	2	1%
Race/Ethnicity										
White, not Hispanic	191	69%	209	69%	226	68%	238	69%	251	70%
Black/African American, not Hispanic	73	26%	81	27%	89	27%	92	27%	95	26%
Hispanic/Latino	6	2%	7	2%	7	2%	7	2%	7	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	6	2%	7	2%	8	2%	8	2%	8	2%
Race/Sex										
White, not Hispanic Males	161	58%	177	58%	192	58%	202	59%	211	58%
White, not Hispanic Females	30	11%	32	11%	34	10%	36	10%	40	11%
Black/African American, not Hispanic Males	49	18%	56	18%	62	19%	65	19%	68	19%
Black/African American, not Hispanic Females	24	9%	25	8%	27	8%	27	8%	27	7%
Hispanic/Latino Males	6	2%	7	2%	7	2%	7	2%	7	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	6	2%	7	2%	8	2%	8	2%	8	2%
Total	276		304		330		345		361	

Notes: Living with a diagnosis of HIV infection by year (2009-2013) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2013.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by year and by transmission category, Stark County

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2009		2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	157	71%	176	72%	194	72%	205	73%	213	73%
Injection drug use (IDU)	2	1%	2	1%	2	1%	2	1%	3	1%
Male-to-male sexual contact and IDU	6	3%	8	3%	8	3%	9	3%	9	3%
Heterosexual contact	11	5%	14	6%	16	6%	16	6%	16	5%
Other/unknown	45	20%	46	19%	48	18%	49	17%	52	18%
Subtotal	221	100%	246	100%	268	100%	281	100%	293	100%
Female adult or adolescent										
Injection drug use	5	10%	5	9%	6	10%	6	10%	6	9%
Heterosexual contact	33	65%	36	67%	39	67%	40	67%	41	64%
Other/unknown	13	25%	13	24%	13	22%	14	23%	17	27%
Subtotal	51	100%	54	100%	58	100%	60	100%	64	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	100%	4	100%	4	100%	4	100%	4	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	4	100%	4	100%	4	100%	4	100%
Total	276		304		330		345		361	

Notes: Living with a diagnosis of HIV infection by year (2009-2013) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2013.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.

Reported persons living with a diagnosis of HIV infection as of June 30, 2014, by year and by exposure category, Stark County

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2009		2010		2011		2012		2013	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	151	55%	169	56%	185	56%	196	57%	204	57%
Injection drug use (IDU) only	4	1%	4	1%	5	2%	5	1%	6	2%
Heterosexual contact only	44	16%	50	16%	55	17%	56	16%	57	16%
MSM & IDU	3	1%	4	1%	4	1%	4	1%	4	1%
IDU & Heterosexual contact	3	1%	3	1%	3	1%	3	1%	3	1%
MSM & Heterosexual contact	6	2%	7	2%	9	3%	9	3%	9	2%
MSM & IDU & Heterosexual contact	3	1%	4	1%	4	1%	5	1%	5	1%
Perinatal exposure	4	1%	4	1%	4	1%	4	1%	4	1%
Other/unknown	58	21%	59	19%	61	18%	63	18%	69	19%
Total	276		304		330		345		361	

Notes: Living with a diagnosis of HIV infection by year (2009-2013) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2013. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2013.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2013.