

## REPORTED CASES OF SELECTED NOTIFIABLE DISEASES BY MONTH OF ONSET, OHIO, 2010

GENERAL INFECTIOUS DISEASES	January		February		March		April		May		June		July	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Amebiasis	4	14%	1	3%	1	3%	6	21%	2	7%	1	3%	2	7%
Botulism	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	33%
Infant*	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	50%
Wound	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Campylobacteriosis	43	4%	50	4%	66	6%	66	6%	109	10%	165	15%	179	16%
Cholera	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	3	100%
Coccidioidomycosis	0	0%	1	6%	3	18%	0	0%	1	6%	0	0%	0	0%
Creutzfeldt-Jakob Disease (CJD)	1	8%	1	8%	1	8%	1	8%	2	17%	1	8%	1	8%
Cryptosporidiosis	29	6%	20	4%	32	7%	30	6%	33	7%	34	7%	71	15%
Cytomegalovirus (CMV), Congenital*	3	11%	0	0%	1	4%	1	4%	1	4%	2	7%	4	14%
<i>Escherichia coli</i> , Shiga Toxin-Producing	2	1%	3	2%	5	4%	18	13%	15	11%	18	13%	29	21%
O157:H7	1	1%	3	4%	4	5%	6	8%	7	9%	11	15%	18	24%
Not O157:H7	1	2%	0	0%	1	2%	12	28%	6	14%	5	12%	10	23%
Unknown Serotype	0	0%	0	0%	0	0%	0	0%	2	10%	2	10%	1	5%
Giardiasis	53	6%	64	7%	69	8%	85	10%	55	6%	68	8%	86	10%
<i>Haemophilus influenzae</i> , Invasive Disease	17	14%	7	6%	5	4%	15	12%	6	5%	18	14%	4	3%
Hemolytic Uremic Syndrome (HUS)	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%	0	0%
Legionellosis	12	5%	18	8%	7	3%	11	5%	24	10%	38	17%	26	11%
Leprosy (Hansen Disease)	0	0%	0	0%	1	100%	0	0%	0	0%	0	0%	0	0%
Listeriosis	1	3%	0	0%	2	7%	1	3%	2	7%	3	10%	6	21%
Meningitis, Aseptic	42	5%	37	5%	48	6%	37	5%	54	7%	49	6%	100	12%
Meningitis, Other Bacterial*	9	11%	3	4%	4	5%	5	6%	6	7%	7	9%	10	12%
Meningococcal Disease	2	6%	5	14%	2	6%	5	14%	2	6%	1	3%	3	9%
Salmonellosis	69	5%	62	5%	73	6%	91	7%	183	14%	164	13%	160	12%
Shigellosis	32	11%	13	4%	13	4%	17	6%	48	16%	36	12%	39	13%
<i>Staphylococcus aureus</i> , Intermediate Resistance to Vancomycin (VISA)	0	0%	0	0%	2	20%	2	20%	2	20%	1	10%	1	10%
Streptococcal Disease, Group A, Invasive	20	8%	20	8%	24	10%	28	11%	22	9%	18	7%	14	6%
Streptococcal Disease, Group B, in Newborn*	2	5%	1	2%	4	10%	5	12%	3	7%	3	7%	4	10%
Streptococcal Toxic Shock Syndrome (STSS)	4	33%	2	17%	1	8%	0	0%	1	8%	1	8%	1	8%
<i>Streptococcus pneumoniae</i> , Invasive Disease	129	11%	119	10%	150	12%	132	11%	119	10%	62	5%	37	3%
Ages < 5 Years*	5	5%	10	10%	13	13%	11	11%	8	8%	8	8%	5	5%
Drug Resistant, Ages 5+ Years*	33	10%	33	10%	37	12%	41	13%	32	10%	17	5%	10	3%
Drug Susceptible, Ages 5+ Years*	91	11%	76	9%	100	12%	80	10%	79	10%	37	5%	22	3%
Toxic Shock Syndrome (TSS)	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	25%
Typhoid Fever	2	22%	0	0%	0	0%	0	0%	1	11%	2	22%	1	11%
Vibriosis	0	0%	0	0%	0	0%	0	0%	1	9%	2	18%	4	36%
<i>Vibrio parahaemolyticus</i> Infection	0	0%	0	0%	0	0%	0	0%	0	0%	2	40%	1	20%
Other (Not Cholera)	0	0%	0	0%	0	0%	0	0%	1	17%	0	0%	3	50%
Yersiniosis	7	17%	5	12%	1	2%	3	7%	1	2%	3	7%	3	7%
<b>SUB-TOTAL</b>	<b>483</b>	<b>7%</b>	<b>432</b>	<b>6%</b>	<b>515</b>	<b>7%</b>	<b>559</b>	<b>8%</b>	<b>693</b>	<b>10%</b>	<b>698</b>	<b>10%</b>	<b>790</b>	<b>11%</b>

### HEPATITIS

Hepatitis A	5	10%	2	4%	0	0%	4	8%	1	2%	2	4%	3	6%
Hepatitis B*	240	13%	148	8%	176	9%	145	8%	182	10%	129	7%	139	7%
Acute*	11	9%	12	10%	12	10%	11	9%	13	11%	17	14%	12	10%
Chronic*	229	13%	135	8%	164	9%	134	8%	169	10%	112	6%	127	7%
Perinatal Infection*	0	0%	1	33%	0	0%	0	0%	0	0%	0	0%	0	0%
Hepatitis C*	595	8%	701	9%	698	9%	597	8%	836	11%	586	8%	708	9%
Acute*	0	0%	2	17%	0	0%	2	17%	0	0%	2	17%	2	17%
Past or Present*	595	8%	699	9%	698	9%	595	8%	836	11%	584	8%	706	9%
Hepatitis E	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
<b>SUB-TOTAL</b>	<b>840</b>	<b>9%</b>	<b>851</b>	<b>9%</b>	<b>874</b>	<b>9%</b>	<b>746</b>	<b>8%</b>	<b>1,019</b>	<b>11%</b>	<b>717</b>	<b>7%</b>	<b>850</b>	<b>9%</b>

N = number of cases reported.

% = percentage of cases occurring in the month for the disease.

\* Please see Technical Notes (pp. 93-97).

## REPORTED CASES OF SELECTED NOTIFIABLE DISEASES BY MONTH OF ONSET, OHIO, 2010

OUTBREAKS*	January		February		March		April		May		June		July	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Community*	2	4%	1	2%	2	4%	4	9%	3	6%	3	6%	1	2%
Foodborne*	8	12%	4	6%	9	13%	7	10%	8	12%	5	7%	4	6%
Healthcare-Associated*	2	3%	7	10%	12	18%	6	9%	4	6%	2	3%	3	4%
Institutional*	5	6%	3	4%	9	11%	9	11%	4	5%	6	7%	1	1%
Waterborne*	0	0%	0	0%	2	20%	0	0%	0	0%	0	0%	1	10%
Zoonotic*	0	0%	0	0%	1	50%	0	0%	1	50%	0	0%	0	0%
<b>SUB-TOTAL</b>	<b>17</b>	<b>6%</b>	<b>15</b>	<b>5%</b>	<b>35</b>	<b>13%</b>	<b>26</b>	<b>9%</b>	<b>20</b>	<b>7%</b>	<b>16</b>	<b>6%</b>	<b>10</b>	<b>4%</b>

### VACCINE-PREVENTABLE

Influenza-Associated Hospitalization*	41	16%	22	8%	15	6%	4	2%	0	0%	1	0%	0	0%
Measles	0	0%	0	0%	0	0%	2	100%	0	0%	0	0%	0	0%
Imported	0	0%	0	0%	0	0%	1	100%	0	0%	0	0%	0	0%
Indigenous	0	0%	0	0%	0	0%	1	100%	0	0%	0	0%	0	0%
Mumps	0	0%	4	15%	1	4%	2	7%	1	4%	0	0%	1	4%
Pertussis	100	5%	54	3%	75	4%	119	6%	138	7%	155	8%	228	12%
Tetanus	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Varicella	121	9%	125	9%	156	12%	150	11%	185	14%	52	4%	34	3%
<b>SUB-TOTAL</b>	<b>262</b>	<b>8%</b>	<b>205</b>	<b>6%</b>	<b>247</b>	<b>7%</b>	<b>277</b>	<b>8%</b>	<b>324</b>	<b>9%</b>	<b>208</b>	<b>6%</b>	<b>263</b>	<b>8%</b>

### ZOOLOSES

Brucellosis	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Dengue	4	25%	1	6%	0	0%	0	0%	0	0%	0	0%	0	0%
Ehrlichiosis/Anaplasmosis	0	0%	0	0%	0	0%	0	0%	0	0%	1	10%	6	60%
<i>Anaplasma phagocytophilum</i> *	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	50%
<i>Ehrlichia chaffeensis</i> *	0	0%	0	0%	0	0%	0	0%	0	0%	1	13%	5	63%
LaCrosse Virus Disease*	0	0%	0	0%	0	0%	0	0%	0	0%	1	4%	4	17%
Lyme Disease	0	0%	1	3%	2	5%	3	8%	3	8%	8	22%	8	22%
Malaria	2	5%	4	9%	2	5%	2	5%	4	9%	7	16%	9	20%
Psittacosis	1	100%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Q Fever	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Rabies, Animal*	2	4%	0	0%	0	0%	2	4%	4	9%	12	26%	4	9%
Rocky Mountain Spotted Fever (RMSF)	0	0%	0	0%	0	0%	2	13%	1	6%	6	38%	5	31%
West Nile Virus Infection*	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	1	20%
<b>SUB-TOTAL</b>	<b>9</b>	<b>4%</b>	<b>6</b>	<b>3%</b>	<b>4</b>	<b>2%</b>	<b>9</b>	<b>4%</b>	<b>12</b>	<b>6%</b>	<b>35</b>	<b>17%</b>	<b>38</b>	<b>19%</b>

<b>GRAND TOTAL</b>	<b>1,611</b>	<b>8%</b>	<b>1,509</b>	<b>7%</b>	<b>1,675</b>	<b>8%</b>	<b>1,617</b>	<b>8%</b>	<b>2,068</b>	<b>10%</b>	<b>1,674</b>	<b>8%</b>	<b>1,951</b>	<b>9%</b>
--------------------	--------------	-----------	--------------	-----------	--------------	-----------	--------------	-----------	--------------	------------	--------------	-----------	--------------	-----------

N = number of cases reported.

% = percentage of cases occurring in the month for the disease.

\* Please see Technical Notes (pp. 93-97).

## REPORTED CASES OF SELECTED NOTIFIABLE DISEASES BY MONTH OF ONSET, OHIO, 2010

GENERAL INFECTIOUS DISEASES	August		September		October		November		December		TOTAL	
	N	%	N	%	N	%	N	%	N	%	N	%
Amebiasis	1	3%	4	14%	0	0%	5	17%	2	7%	29	100%
Botulism	0	0%	1	33%	0	0%	0	0%	1	33%	3	100%
Infant*	0	0%	0	0%	0	0%	0	0%	1	50%	2	100%
Wound	0	0%	1	100%	0	0%	0	0%	0	0%	1	100%
Campylobacteriosis	143	13%	107	10%	77	7%	62	6%	57	5%	1,124	100%
Cholera	0	0%	0	0%	0	0%	0	0%	0	0%	3	100%
Coccidioidomycosis	3	18%	1	6%	4	24%	1	6%	3	18%	17	100%
Creutzfeldt-Jakob Disease (CJD)	2	17%	0	0%	0	0%	2	17%	0	0%	12	100%
Cryptosporidiosis	71	15%	71	15%	34	7%	19	4%	33	7%	477	100%
Cytomegalovirus (CMV), Congenital*	1	4%	1	4%	7	25%	2	7%	5	18%	28	100%
<i>Escherichia coli</i> , Shiga Toxin-Producing	24	17%	4	3%	10	7%	5	4%	5	4%	138	100%
O157:H7	12	16%	3	4%	4	5%	5	7%	1	1%	75	100%
Not O157:H7	7	16%	0	0%	1	2%	0	0%	0	0%	43	100%
Unknown Serotype	5	25%	1	5%	5	25%	0	0%	4	20%	20	100%
Giardiasis	80	9%	80	9%	84	10%	65	8%	74	9%	863	100%
<i>Haemophilus influenzae</i> , Invasive Disease	9	7%	9	7%	9	7%	14	11%	12	10%	125	100%
Hemolytic Uremic Syndrome (HUS)	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Legionellosis	27	12%	16	7%	21	9%	13	6%	17	7%	230	100%
Leprosy (Hansen Disease)	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Listeriosis	6	21%	5	17%	2	7%	1	3%	0	0%	29	100%
Meningitis, Aseptic	156	19%	99	12%	84	10%	43	5%	61	8%	810	100%
Meningitis, Other Bacterial*	7	9%	9	11%	4	5%	13	16%	5	6%	82	100%
Meningococcal Disease	2	6%	3	9%	3	9%	2	6%	5	14%	35	100%
Salmonellosis	140	11%	108	8%	109	8%	74	6%	76	6%	1,309	100%
Shigellosis	27	9%	14	5%	22	7%	22	7%	21	7%	304	100%
<i>Staphylococcus aureus</i> , Intermediate Resistance to Vancomycin (VISA)	0	0%	1	10%	1	10%	0	0%	0	0%	10	100%
Streptococcal Disease, Group A, Invasive	16	6%	14	6%	13	5%	29	12%	30	12%	248	100%
Streptococcal Disease, Group B, in Newborn*	3	7%	4	10%	4	10%	5	12%	3	7%	41	100%
Streptococcal Toxic Shock Syndrome (STSS)	1	8%	0	0%	0	0%	0	0%	1	8%	12	100%
<i>Streptococcus pneumoniae</i> , Invasive Disease	34	3%	60	5%	109	9%	133	11%	136	11%	1,220	100%
Ages < 5 Years*	4	4%	9	9%	9	9%	5	5%	10	10%	97	100%
Drug Resistant, Ages 5+ Years*	9	3%	14	4%	25	8%	35	11%	34	11%	320	100%
Drug Susceptible, Ages 5+ Years*	21	3%	37	5%	75	9%	93	12%	92	11%	803	100%
Toxic Shock Syndrome (TSS)	1	25%	0	0%	1	25%	0	0%	1	25%	4	100%
Typhoid Fever	1	11%	1	11%	1	11%	0	0%	0	0%	9	100%
Vibriosis	2	18%	1	9%	0	0%	1	9%	0	0%	11	100%
<i>Vibrio parahaemolyticus</i> Infection	1	20%	1	20%	0	0%	0	0%	0	0%	5	100%
Other (Not Cholera)	1	17%	0	0%	0	0%	1	17%	0	0%	6	100%
Yersiniosis	4	10%	1	2%	5	12%	3	7%	6	14%	42	100%
<b>SUB-TOTAL</b>	<b>761</b>	<b>11%</b>	<b>614</b>	<b>9%</b>	<b>604</b>	<b>8%</b>	<b>514</b>	<b>7%</b>	<b>554</b>	<b>8%</b>	<b>7,217</b>	<b>100%</b>

### HEPATITIS

Hepatitis A	11	22%	10	20%	5	10%	2	4%	4	8%	49	100%
Hepatitis B*	130	7%	123	6%	194	10%	143	8%	154	8%	1,903	100%
Acute*	13	11%	5	4%	5	4%	6	5%	6	5%	123	100%
Chronic*	116	7%	117	7%	189	11%	137	8%	148	8%	1,777	100%
Perinatal Infection*	1	33%	1	33%	0	0%	0	0%	0	0%	3	100%
Hepatitis C*	583	8%	503	7%	694	9%	587	8%	640	8%	7,728	100%
Acute*	1	8%	1	8%	0	0%	0	0%	2	17%	12	100%
Past or Present*	582	8%	502	7%	694	9%	587	8%	638	8%	7,716	100%
Hepatitis E	0	100%	1	100%	0	100%	0	100%	0	100%	1	100%
<b>SUB-TOTAL</b>	<b>724</b>	<b>7%</b>	<b>637</b>	<b>7%</b>	<b>893</b>	<b>9%</b>	<b>732</b>	<b>8%</b>	<b>798</b>	<b>8%</b>	<b>9,681</b>	<b>100%</b>

N = number of cases reported.

% = percentage of cases occurring in the month for the disease.

\* Please see Technical Notes (pp. 93-97).

## REPORTED CASES OF SELECTED NOTIFIABLE DISEASES BY MONTH OF ONSET, OHIO, 2010

OUTBREAKS*	August		September		October		November		December		TOTAL	
	N	%	N	%	N	%	N	%	N	%	N	%
Community*	2	4%	9	19%	9	19%	7	15%	4	9%	47	100%
Foodborne*	1	1%	8	12%	4	6%	5	7%	6	9%	69	100%
Healthcare-Associated*	3	4%	4	6%	4	6%	8	12%	13	19%	68	100%
Institutional*	1	1%	3	4%	9	11%	13	16%	19	23%	82	100%
Waterborne*	0	0%	1	10%	0	0%	0	0%	6	60%	10	100%
Zoonotic*	0	0%	0	0%	0	0%	0	0%	0	0%	2	100%
<b>SUB-TOTAL</b>	<b>7</b>	<b>3%</b>	<b>25</b>	<b>9%</b>	<b>26</b>	<b>9%</b>	<b>33</b>	<b>12%</b>	<b>48</b>	<b>17%</b>	<b>278</b>	<b>100%</b>

VACCINE-PREVENTABLE												
Influenza-Associated Hospitalization*	0	0%	4	2%	14	5%	30	12%	128	49%	259	100%
Measles	0	0%	0	0%	0	0%	0	0%	0	0%	2	100%
Imported	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Indigenous	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Mumps	1	4%	5	19%	8	30%	3	11%	1	4%	27	100%
Pertussis	161	9%	235	13%	229	12%	222	12%	142	8%	1,858	100%
Tetanus	0	0%	0	0%	1	100%	0	0%	0	0%	1	100%
Varicella	43	3%	107	8%	157	12%	117	9%	90	7%	1,337	100%
<b>SUB-TOTAL</b>	<b>205</b>	<b>6%</b>	<b>351</b>	<b>10%</b>	<b>409</b>	<b>12%</b>	<b>372</b>	<b>11%</b>	<b>361</b>	<b>10%</b>	<b>3,484</b>	<b>100%</b>

ZOOZOSES												
Brucellosis	0	0%	0	0%	0	0%	1	100%	0	0%	1	100%
Dengue	8	50%	1	6%	2	13%	0	0%	0	0%	16	100%
Ehrlichiosis/Anaplasmosis	2	20%	0	0%	0	0%	1	10%	0	0%	10	100%
<i>Anaplasma phagocytophilum</i> *	1	50%	0	0%	0	0%	0	0%	0	0%	2	100%
<i>Ehrlichia chaffeensis</i> *	1	13%	0	0%	0	0%	1	13%	0	0%	8	100%
LaCrosse Virus Disease*	14	58%	5	21%	0	0%	0	0%	0	0%	24	100%
Lyme Disease	5	14%	1	3%	3	8%	2	5%	1	3%	37	100%
Malaria	1	2%	3	7%	3	7%	3	7%	4	9%	44	100%
Psittacosis	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Q Fever	0	0%	0	0%	0	0%	0	0%	0	0%	1	100%
Rabies, Animal*	18	38%	2	4%	1	2%	1	2%	1	2%	47	100%
Rocky Mountain Spotted Fever (RMSF)	1	6%	0	0%	1	6%	0	0%	0	0%	16	100%
West Nile Virus Infection*	1	20%	2	40%	1	20%	0	0%	0	0%	5	100%
<b>SUB-TOTAL</b>	<b>50</b>	<b>25%</b>	<b>14</b>	<b>7%</b>	<b>11</b>	<b>5%</b>	<b>8</b>	<b>4%</b>	<b>6</b>	<b>3%</b>	<b>202</b>	<b>100%</b>

<b>GRAND TOTAL</b>	<b>1,747</b>	<b>8%</b>	<b>1,641</b>	<b>8%</b>	<b>1,943</b>	<b>9%</b>	<b>1,659</b>	<b>8%</b>	<b>1,767</b>	<b>8%</b>	<b>20,862</b>	<b>100%</b>
--------------------	--------------	-----------	--------------	-----------	--------------	-----------	--------------	-----------	--------------	-----------	---------------	-------------

N = number of cases reported.

% = percentage of cases occurring in the month for the disease.

\* Please see Technical Notes (pp. 93-97).