

Section 4: Ryan White Part A and Part B Regional Summary Data

Section 4 presents summary data for Ryan White Part A Cleveland and Columbus Planning Regions and the 11 Ryan White Part B Consortia Areas. Data presented include HIV/AIDS surveillance data on reported persons living with a diagnosis of HIV infection, reported diagnoses of HIV infection and reported cases of AIDS.

Ryan White Part B Consortia Areas



Table 142. Breakdown of Ryan White Part B Consortia and Part A Planning Regions

Breakdown of Ryan White Part B and Part A Planning Regions			
<p>1 Cleveland Ashtabula Lake Cuyahoga Lorain Geauga</p> <p>2 Columbus Delaware Madison Fairfield Pickaway Franklin Union Licking</p> <p>3 Cincinnati Adams Clinton Brown Hamilton Butler Highland Clermont Warren</p> <p>4 Dayton Clark Miami Darke Montgomery Greene Preble</p> <p>5 Toledo Defiance Ottawa Fulton Sandusky Henry Williams Lucas Wood</p> <p>6 Akron Medina Portage Summit</p> <p>7 Youngstown Columbiana Jefferson Mahoning Trumbull</p> <p>8 Canton Carroll Stark Harrison Tuscarawas Holmes Wayne</p>	<p>9A Lima Allen Mercer Auglaize Paulding Champaign Putnam Hancock Shelby Hardin Van Wert Logan</p> <p>9B Mansfield Ashland Marion Crawford Morrow Erie Richland Huron Seneca Knox Wyandot</p> <p>9C Southeast Athens Monroe Belmont Morgan Coshocton Muskingum Fayette Noble Gallia Perry Guernsey Pike Hocking Ross Jackson Scioto Lawrence Vinton Meigs Washington</p>		
Part A Planning Regions			
Cleveland			
Ashtabula			
Cuyahoga			
Geauga			
Lake			
Lorain			
Medina			
Columbus			
Delaware Madison			
Fairfield Morrow			
Franklin Pickaway			
Licking Union			

HIV/AIDS Data by Ryan White Part A Cleveland Planning Region and Ryan White Part B Consortia Areas

The following tables present HIV/AIDS data for Ryan White Part A Cleveland and Columbus Planning Region and for the 11 Ryan White Part B Consortia Areas. There are eight data tables for each Ryan White Part:

- Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics
- Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category
- Reported persons living with a diagnosis of HIV infection by year and selected characteristics, 2007-2011
- Reported persons living with a diagnosis of HIV infection by year and transmission category, 2007-2011
- Reported diagnoses of HIV infection by year of diagnosis and cumulative diagnoses by selected characteristics, 2007-2011
- Reported diagnoses of HIV infection by year of diagnosis and cumulative diagnoses by selected characteristics, 2007-2011
- Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, 2007-2011
- Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, 2007-2011

When viewing these tables, it is important to take into account the limitations of the data. These data represent confidential reports of HIV infection and AIDS diagnoses; not all persons with HIV infection. The HIV diagnosis data represent the earliest date of diagnosis reported to the HIV/AIDS Surveillance Program. The earliest date reported may not be the earliest date an individual became aware of their HIV infection. Individuals may have previously tested anonymously or were diagnosed out-of-state prior to being confidentially tested and reported to Ohio.

Ryan White Part A Cleveland Planning Region

Table 143. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Cleveland Part A

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	317.7	3,322	77%	1,658	77%	1,664	78%
Females	86.0	966	23%	500	23%	466	22%
Age at diagnosis (yr)							
<13	4.1	14	<1%	9	<1%	5	<1%
13-14	10.3	6	<1%	4	<1%	2	<1%
15-19	16.8	25	1%	19	1%	6	<1%
20-24	122.2	154	4%	121	6%	33	2%
25-29	201.7	259	6%	192	9%	67	3%
30-34	253.5	322	8%	215	10%	107	5%
35-39	269.7	338	8%	202	9%	136	6%
40-44	447.2	649	15%	327	15%	322	15%
45-49	493.9	789	18%	353	16%	436	20%
50-54	434.7	762	18%	307	14%	455	21%
55-64	265.9	790	18%	332	15%	458	22%
65+	53.8	180	4%	77	4%	103	5%
Race/Ethnicity							
White, not Hispanic	97.9	1,540	36%	762	35%	778	37%
Black/African American, not Hispanic	552.3	2,274	53%	1,156	54%	1,118	52%
Hispanic/Latino	355.4	369	9%	170	8%	199	9%
Asian/Pacific Islander	35.4	15	<1%	11	1%	4	<1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	86	2%	57	3%	29	1%
Race/Sex							
White, not Hispanic Males	175.5	1,343	31%	654	30%	689	32%
White, not Hispanic Females	24.4	197	5%	108	5%	89	4%
Black/African American, not Hispanic Males	869.8	1,651	39%	837	39%	814	38%
Black/African American, not Hispanic Females	280.8	623	15%	319	15%	304	14%
Hispanic/Latino Males	474.7	247	6%	114	5%	133	6%
Hispanic/Latina Females	235.5	122	3%	56	3%	66	3%
Asian/Pacific Islander Males	54.6	11	<1%	8	<1%	3	<1%
Asian/Pacific Islander Females	*	4	<1%	3	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	86	2%	57	3%	29	1%
Total	197.6	4,288		2,158		2,130	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 144. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Cleveland Part A

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,174	66%	1,045	64%	1,129	68%
Injection drug use (IDU)	190	6%	69	4%	121	7%
Male-to-male sexual contact and IDU	150	5%	45	3%	105	6%
Heterosexual contact	241	7%	115	7%	126	8%
Other/unknown	538	16%	367	22%	171	10%
Subtotal	3,293	100%	1,641	100%	1,652	100%
Female adult or adolescent						
Injection drug use	122	13%	39	8%	83	18%
Heterosexual contact	577	62%	279	58%	298	66%
Other/unknown	238	25%	166	34%	72	16%
Subtotal	937	100%	484	100%	453	100%
Child (<13 yrs at diagnosis)						
Perinatal	50	86%	27	82%	23	92%
Other/unknown	8	14%	6	18%	2	8%
Subtotal	58	100%	33	100%	25	100%
Total	4,288		2,158		2,130	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 145. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Cleveland Part A, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	2,630	77%	2,816	77%	2,983	77%	3,139	77%	3,322	77%
Females	770	23%	829	23%	886	23%	920	23%	966	23%
Age at diagnosis (yr)										
<13	11	<1%	12	<1%	14	<1%	14	<1%	14	<1%
13-14	5	<1%	6	<1%	6	<1%	6	<1%	6	<1%
15-19	18	1%	18	<1%	18	<1%	21	1%	25	1%
20-24	30	1%	46	1%	70	2%	110	3%	154	4%
25-29	139	4%	176	5%	204	5%	230	6%	259	6%
30-34	192	6%	227	6%	263	7%	286	7%	322	8%
35-39	249	7%	278	8%	298	8%	321	8%	338	8%
40-44	531	16%	567	16%	606	16%	623	15%	649	15%
45-49	693	20%	720	20%	746	19%	765	19%	789	18%
50-54	667	20%	701	19%	720	19%	739	18%	762	18%
55-64	696	20%	721	20%	748	19%	767	19%	790	18%
65+	169	5%	173	5%	176	5%	177	4%	180	4%
Race/Ethnicity										
White, not Hispanic	1,257	37%	1,341	37%	1,413	37%	1,469	36%	1,540	36%
Black/African American, not Hispanic	1,753	52%	1,889	52%	2,019	52%	2,133	53%	2,274	53%
Hispanic/Latino	300	9%	323	9%	339	9%	355	9%	369	9%
Asian/Pacific Islander	13	<1%	13	<1%	14	<1%	15	<1%	15	<1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	73	2%	75	2%	80	2%	83	2%	86	2%
Race/Sex										
White, not Hispanic Males	1,095	32%	1,170	32%	1,232	32%	1,283	32%	1,343	31%
White, not Hispanic Females	162	5%	171	5%	181	5%	186	5%	197	5%
Black/African American, not Hispanic Males	1,265	37%	1,360	37%	1,451	38%	1,541	38%	1,651	39%
Black/African American, not Hispanic Females	488	14%	529	15%	568	15%	592	15%	623	15%
Hispanic/Latino Males	200	6%	214	6%	225	6%	236	6%	247	6%
Hispanic/Latina Females	100	3%	109	3%	114	3%	119	3%	122	3%
Asian/Pacific Islander Males	10	<1%	10	<1%	10	<1%	11	<1%	11	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	4	<1%	4	<1%	4	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	73	2%	75	2%	80	2%	83	2%	86	2%
Total	3,400		3,645		3,869		4,059		4,288	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 146. Reported persons living with a diagnosis of HIV infection by year and transmission category, Cleveland Part A, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,699	65%	1,833	66%	1,949	66%	2,049	66%	2,174	66%
Injection drug use (IDU)	177	7%	181	6%	187	6%	187	6%	190	6%
Male-to-male sexual contact and IDU	133	5%	142	5%	146	5%	148	5%	150	5%
Heterosexual contact	198	8%	214	8%	232	8%	235	8%	241	7%
Other/unknown	396	15%	419	15%	440	15%	491	16%	538	16%
Subtotal	2,603	100%	2,789	100%	2,954	100%	3,110	100%	3,293	100%
Female adult or adolescent										
Injection drug use	108	15%	111	14%	114	13%	118	13%	122	13%
Heterosexual contact	476	64%	518	65%	552	64%	561	63%	577	62%
Other/unknown	159	21%	171	21%	191	22%	212	24%	238	25%
Subtotal	743	100%	800	100%	857	100%	891	100%	937	100%
Child (<13 yrs at diagnosis)										
Perinatal	47	87%	48	86%	50	86%	50	86%	50	86%
Other/unknown	7	13%	8	14%	8	14%	8	14%	8	14%
Subtotal	54	100%	56	100%	58	100%	58	100%	58	100%
Total	3,400		3,645		3,869		4,059		4,288	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 147. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Cleveland Part A, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	19.0	199	83%	19.1	200	76%	16.8	176	75%	15.3	160	82%	17.8	186	80%
Females	3.6	40	17%	5.7	64	24%	5.3	60	25%	3.1	35	18%	4.1	46	20%
Age at diagnosis (yr)															
<13	*	1	<1%	*	2	1%	*	2	1%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
15-19	8.0	12	5%	8.0	12	5%	7.4	11	5%	12.7	19	10%	9.4	14	6%
20-24	32.5	41	17%	29.4	37	14%	23.8	30	13%	28.6	36	18%	33.3	42	18%
25-29	19.5	25	10%	28.8	37	14%	30.4	39	17%	20.2	26	13%	25.7	33	14%
30-34	17.3	22	9%	23.6	30	11%	14.2	18	8%	20.5	26	13%	23.6	30	13%
35-39	30.3	38	16%	27.9	35	13%	30.3	38	16%	16.8	21	11%	12.0	15	6%
40-44	24.1	35	15%	20.7	30	11%	17.9	26	11%	11.0	16	8%	20.0	29	13%
45-49	18.8	30	13%	21.3	34	13%	19.4	31	13%	13.8	22	11%	18.8	30	13%
50-54	10.3	18	8%	13.7	24	9%	11.4	20	8%	8.0	14	7%	10.8	19	8%
55-64	3.4	10	4%	7.1	21	8%	5.7	17	7%	4.7	14	7%	6.1	18	8%
65+	2.1	7	3%	*	2	1%	*	4	2%	*	-	-	*	2	1%
Race/Ethnicity															
White, not Hispanic	5.2	82	34%	5.8	91	34%	5.0	78	33%	3.8	60	31%	4.6	72	31%
Black/African American, not Hispanic	32.8	135	56%	35.5	146	55%	33.0	136	58%	27.9	115	59%	34.7	143	62%
Hispanic/Latino	19.3	20	8%	23.1	24	9%	15.4	16	7%	15.4	16	8%	13.5	14	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	1	<1%	*	1	1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	3	1%	*	5	2%	*	3	2%	*	3	1%
Race/Sex															
White, not Hispanic Males	9.8	75	31%	10.6	81	31%	8.8	67	28%	7.1	54	28%	8.0	61	26%
White, not Hispanic Females	0.9	7	3%	1.2	10	4%	1.4	11	5%	0.7	6	3%	1.4	11	5%
Black/African American, not Hispanic Males	56.4	107	45%	54.3	103	39%	50.0	95	40%	47.9	91	47%	59.0	112	48%
Black/African American, not Hispanic Females	12.6	28	12%	19.4	43	16%	18.5	41	17%	10.8	24	12%	14.0	31	13%
Hispanic/Latino Males	28.8	15	6%	26.9	14	5%	21.1	11	5%	21.1	11	6%	21.1	11	5%
Hispanic/Latina Females	9.7	5	2%	19.3	10	4%	9.7	5	2%	9.7	5	3%	*	3	1%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	3	1%	*	5	2%	*	3	2%	*	3	1%
Total	11.0	239		12.2	264		10.9	236		9.0	195		10.7	232	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 148. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Cleveland Part A, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	140	71%	144	72%	114	66%	103	64%	127	68%
Injection drug use (IDU)	4	2%	3	2%	7	4%	-	-	3	2%
Male-to-male sexual contact and IDU	4	2%	8	4%	5	3%	2	1%	2	1%
Heterosexual contact	21	11%	17	9%	18	10%	3	2%	6	3%
Other/unknown	29	15%	28	14%	30	17%	52	33%	48	26%
Subtotal	198	100%	200	100%	174	100%	160	100%	186	100%
Female adult or adolescent										
Injection drug use	3	8%	4	6%	3	5%	4	11%	4	9%
Heterosexual contact	28	70%	45	73%	34	57%	9	26%	16	35%
Other/unknown	9	23%	13	21%	23	38%	22	63%	26	57%
Subtotal	40	100%	62	100%	60	100%	35	100%	46	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	1	50%	2	100%	-	-	-	-
Other/unknown	-	-	1	50%	-	-	-	-	-	-
Subtotal	1	100%	2	100%	2	100%	-	-	-	-
Total	239		264		236		195		232	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 149. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Cleveland Part A, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	83	78%	86	75%	89	72%	73	74%	76	78%	3,812	82%
Females	23	22%	28	25%	34	28%	26	26%	22	22%	830	18%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	48	1%
13-14	1	1%	-	-	-	-	-	-	-	-	3	<1%
15-19	4	4%	-	-	3	2%	3	3%	2	2%	36	1%
20-24	11	10%	9	8%	8	7%	12	12%	6	6%	220	5%
25-29	7	7%	13	11%	8	7%	11	11%	12	12%	589	13%
30-34	8	8%	15	13%	13	11%	12	12%	10	10%	911	20%
35-39	18	17%	17	15%	24	20%	12	12%	5	5%	1,014	22%
40-44	18	17%	22	19%	17	14%	13	13%	19	19%	747	16%
45-49	19	18%	17	15%	23	19%	19	19%	24	24%	526	11%
50-54	12	11%	9	8%	12	10%	11	11%	9	9%	255	5%
55-64	2	2%	11	10%	14	11%	6	6%	9	9%	232	5%
65+	6	6%	1	1%	1	1%	-	-	2	2%	60	1%
Race/Ethnicity												
White, not Hispanic	45	42%	35	31%	40	33%	29	29%	27	28%	1,879	40%
Black/African American, not Hispanic	51	48%	60	53%	69	56%	60	61%	66	67%	2,339	50%
Hispanic/Latino	10	9%	17	15%	11	9%	9	9%	4	4%	376	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	8	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	6	<1%
Unknown	-	-	2	2%	3	2%	1	1%	1	1%	34	1%
Race/Sex												
White, not Hispanic Males	38	36%	32	28%	35	28%	24	24%	23	23%	1,722	37%
White, not Hispanic Females	7	7%	3	3%	5	4%	5	5%	4	4%	157	3%
Black/African American, not Hispanic Males	38	36%	45	39%	46	37%	43	43%	48	49%	1,795	39%
Black/African American, not Hispanic Females	13	12%	15	13%	23	19%	17	17%	18	18%	544	12%
Hispanic/Latino Males	7	7%	8	7%	6	5%	5	5%	4	4%	260	6%
Hispanic/Latina Females	3	3%	9	8%	5	4%	4	4%	-	-	116	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	7	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	3	<1%
Unknown	-	-	2	2%	3	2%	1	1%	1	1%	34	1%
Total	106		114		123		99		98		4,642	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Cleveland Planning Region

Table 150. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Cleveland Part A, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	57	69%	58	67%	54	61%	48	66%	44	59%	2,571	68%
Injection drug use (IDU)	3	4%	2	2%	6	7%	2	3%	1	1%	381	10%
Male-to-male sexual contact and IDU	2	2%	7	8%	6	7%	3	4%	2	3%	282	7%
Heterosexual contact	11	13%	11	13%	12	13%	3	4%	3	4%	211	6%
Other/unknown	10	12%	8	9%	11	12%	17	23%	25	33%	343	9%
Subtotal	83	100%	86	100%	89	100%	73	100%	75	100%	3,788	100%
Female adult or adolescent												
Injection drug use	6	27%	2	7%	4	12%	5	21%	1	5%	193	24%
Heterosexual contact	14	64%	22	79%	25	74%	9	38%	9	41%	471	59%
Other/unknown	2	9%	4	14%	5	15%	10	42%	12	55%	132	17%
Subtotal	22	100%	28	100%	34	100%	24	100%	22	100%	796	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	-	-	2	100%	1	100%	50	86%
Other/unknown	-	-	-	-	-	-	-	-	-	-	8	14%
Subtotal	1	100%	-	-	-	-	2	100%	1	100%	58	100%
Total	106		114		123		99		98		4,642	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 151. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Cleveland Part A

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	375.1	3,423	82%	1,918	81%	1,505	83%
Females	80.1	758	18%	447	19%	311	17%
Age at diagnosis (yr)							
<13	3.9	13	<1%	6	<1%	7	<1%
13-14	*	-	-	-	-	-	-
15-19	18.2	23	1%	17	1%	6	<1%
20-24	144.5	200	5%	154	7%	46	3%
25-29	226.1	318	8%	231	10%	87	5%
30-34	291.9	395	9%	263	11%	132	7%
35-39	349.8	441	11%	250	11%	191	11%
40-44	479.4	637	15%	350	15%	287	16%
45-49	619.1	817	20%	424	18%	393	22%
50-54	459.0	610	15%	300	13%	310	17%
55-64	285.3	607	15%	304	13%	303	17%
65+	60.4	121	3%	67	3%	54	3%
Race/Ethnicity							
White, not Hispanic	168.9	2,379	57%	1,363	58%	1,016	56%
Black/African American, not Hispanic	552.2	1,522	36%	848	36%	674	37%
Hispanic/Latino	215.6	150	4%	74	3%	76	4%
Asian/Pacific Islander	36.2	22	1%	13	1%	9	<1%
American Indian/Alaska Native	*	4	<1%	3	<1%	1	<1%
Unknown	*	105	3%	65	3%	40	2%
Race/Sex							
White, not Hispanic Males	310.0	2,145	51%	1,216	51%	929	51%
White, not Hispanic Females	32.7	234	6%	147	6%	87	5%
Black/African American, not Hispanic Males	792.3	1,049	25%	575	24%	474	26%
Black/African American, not Hispanic Females	330.3	473	11%	273	12%	200	11%
Hispanic/Latino Males	329.2	121	3%	61	3%	60	3%
Hispanic/Latina Females	88.3	29	1%	13	1%	16	1%
Asian/Pacific Islander Males	57.2	17	<1%	10	<1%	7	<1%
Asian/Pacific Islander Females	16.1	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native Males	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	105	3%	65	3%	40	2%
Total	225.0	4,182		2,366		1,816	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 152. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Cleveland Part A

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,319	68%	1,292	68%	1,027	69%
Injection drug use (IDU)	75	2%	21	1%	54	4%
Male-to-male sexual contact and IDU	93	3%	40	2%	53	4%
Heterosexual contact	219	6%	84	4%	135	9%
Other/unknown	701	21%	471	25%	230	15%
Subtotal	3,407	100%	1,908	100%	1,499	100%
Female adult or adolescent						
Injection drug use	61	8%	28	6%	33	11%
Heterosexual contact	396	54%	214	49%	182	60%
Other/unknown	282	38%	196	45%	86	29%
Subtotal	739	100%	438	100%	301	100%
Child (<13 yrs at diagnosis)						
Perinatal	29	83%	16	84%	13	81%
Other/unknown	6	17%	3	16%	3	19%
Subtotal	35	100%	19	100%	16	100%
Total	4,182		2,366		1,816	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 153. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Cleveland Part A, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	2,487	81%	2,713	81%	2,942	81%	3,177	82%	3,423	82%
Females	571	19%	616	18%	671	19%	709	18%	758	18%
Age at diagnosis (yr)										
<13	9	<1%	11	<1%	12	<1%	12	<1%	13	<1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	10	<1%	10	<1%	13	<1%	15	<1%	23	1%
20-24	37	1%	56	2%	102	3%	148	4%	200	5%
25-29	117	4%	165	5%	220	6%	268	7%	318	8%
30-34	225	7%	262	8%	305	8%	348	9%	395	9%
35-39	291	10%	328	10%	364	10%	400	10%	441	11%
40-44	491	16%	529	16%	570	16%	603	16%	637	15%
45-49	706	23%	741	22%	764	21%	792	20%	817	20%
50-54	533	17%	558	17%	573	16%	591	15%	610	15%
55-64	537	18%	562	17%	577	16%	592	15%	607	15%
65+	103	3%	108	3%	114	3%	118	3%	121	3%
Race/Ethnicity										
White, not Hispanic	1,803	59%	1,949	59%	2,095	58%	2,234	57%	2,379	57%
Black/African American, not Hispanic	1,064	35%	1,170	35%	1,289	36%	1,401	36%	1,522	36%
Hispanic/Latino	100	3%	109	3%	119	3%	132	3%	150	4%
Asian/Pacific Islander	14	<1%	16	<1%	18	<1%	20	1%	22	1%
American Indian/Alaska Native	3	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	75	2%	82	2%	89	2%	96	2%	105	3%
Race/Sex										
White, not Hispanic Males	1,624	53%	1,756	53%	1,887	52%	2,014	52%	2,145	51%
White, not Hispanic Females	179	6%	193	6%	208	6%	220	6%	234	6%
Black/African American, not Hispanic Males	710	23%	787	24%	870	24%	959	25%	1,049	25%
Black/African American, not Hispanic Females	354	12%	383	12%	419	12%	442	11%	473	11%
Hispanic/Latino Males	77	3%	86	3%	95	3%	107	3%	121	3%
Hispanic/Latina Females	23	1%	23	1%	24	1%	25	1%	29	1%
Asian/Pacific Islander Males	11	<1%	13	<1%	14	<1%	15	<1%	17	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	4	<1%	5	<1%	5	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	75	2%	82	2%	89	2%	96	2%	105	3%
Total	3,059		3,330		3,614		3,887		4,182	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 154. Reported persons living with a diagnosis of HIV infection by year and transmission category, Cleveland Part A, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,659	67%	1,836	68%	2,006	69%	2,163	68%	2,319	68%
Injection drug use (IDU)	68	3%	69	3%	70	2%	70	2%	75	2%
Male-to-male sexual contact and IDU	66	3%	73	3%	80	3%	88	3%	93	3%
Heterosexual contact	161	7%	187	7%	205	7%	215	7%	219	6%
Other/unknown	517	21%	532	20%	565	19%	625	20%	701	21%
Subtotal	2,471	100%	2,697	100%	2,926	100%	3,161	100%	3,407	100%
Female adult or adolescent										
Injection drug use	54	10%	57	10%	58	9%	61	9%	61	8%
Heterosexual contact	303	54%	329	55%	365	56%	380	55%	396	54%
Other/unknown	200	36%	214	36%	230	35%	250	36%	282	38%
Subtotal	557	100%	600	100%	653	100%	691	100%	739	100%
Child (<13 yrs at diagnosis)										
Perinatal	24	80%	26	81%	28	82%	28	82%	29	83%
Other/unknown	6	20%	6	19%	6	18%	6	18%	6	17%
Subtotal	30	100%	32	100%	34	100%	34	100%	35	100%
Total	3,059		3,330		3,614		3,887		4,182	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 155. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Cleveland Part A, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	27.0	246	83%	26.8	245	84%	26.4	241	81%	26.8	245	86%	27.4	250	83%
Females	5.3	50	17%	4.9	46	16%	6.0	57	19%	4.1	39	14%	5.4	51	17%
Age at diagnosis (yr)															
<13	*	3	1%	*	2	1%	*	2	1%	*	-	-	*	1	<1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	10.3	13	4%	10.3	13	4%	22.2	28	9%	10.3	13	5%	12.7	16	5%
20-24	29.6	41	14%	35.4	49	17%	37.6	52	17%	43.4	60	21%	40.5	56	19%
25-29	41.2	58	20%	26.3	37	13%	40.5	57	19%	30.6	43	15%	32.7	46	15%
30-34	19.2	26	9%	32.5	44	15%	25.9	35	12%	29.6	40	14%	34.0	46	15%
35-39	38.9	49	17%	30.9	39	13%	32.5	41	14%	29.3	37	13%	34.9	44	15%
40-44	40.6	54	18%	31.6	42	14%	27.1	36	12%	27.1	36	13%	26.3	35	12%
45-49	17.4	23	8%	22.0	29	10%	13.6	18	6%	19.7	26	9%	16.7	22	7%
50-54	14.3	19	6%	18.8	25	9%	10.5	14	5%	9.0	12	4%	15.0	20	7%
55-64	4.2	9	3%	2.8	6	2%	5.6	12	4%	7.1	15	5%	5.6	12	4%
65+	*	1	<1%	2.5	5	2%	*	3	1%	*	2	1%	*	3	1%
Race/Ethnicity															
White, not Hispanic	11.4	160	54%	11.1	156	54%	10.9	154	52%	10.3	145	51%	10.6	149	50%
Black/African American, not Hispanic	40.3	111	38%	41.7	115	40%	45.4	125	42%	42.4	117	41%	44.6	123	41%
Hispanic/Latino	20.1	14	5%	14.4	10	3%	14.4	10	3%	18.7	13	5%	25.9	18	6%
Asian/Pacific Islander	*	3	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
American Indian/Alaska Native	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	8	3%	*	7	2%	*	7	2%	*	7	2%	*	9	3%
Race/Sex															
White, not Hispanic Males	21.0	145	49%	20.5	142	49%	20.1	139	47%	19.2	133	47%	19.4	134	45%
White, not Hispanic Females	2.1	15	5%	2.0	14	5%	2.1	15	5%	1.7	12	4%	2.1	15	5%
Black/African American, not Hispanic Males	59.7	79	27%	65.0	86	30%	65.7	87	29%	70.2	93	33%	68.7	91	30%
Black/African American, not Hispanic Females	22.3	32	11%	20.2	29	10%	26.5	38	13%	16.8	24	8%	22.3	32	11%
Hispanic/Latino Males	35.4	13	4%	24.5	9	3%	24.5	9	3%	32.7	12	4%	38.1	14	5%
Hispanic/Latina Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	4	1%
Asian/Pacific Islander Males	*	3	1%	*	2	1%	*	1	<1%	*	1	<1%	*	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	1	<1%	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	8	3%	*	7	2%	*	7	2%	*	7	2%	*	9	3%
Total	15.9	296		15.7	291		16.0	298		15.3	284		16.2	301	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 156. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Cleveland Part A, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	188	77%	191	78%	178	74%	162	66%	157	63%
Injection drug use (IDU)	6	2%	2	1%	1	<1%	-	-	5	2%
Male-to-male sexual contact and IDU	8	3%	9	4%	7	3%	8	3%	5	2%
Heterosexual contact	12	5%	27	11%	18	7%	10	4%	4	2%
Other/unknown	31	13%	16	7%	37	15%	65	27%	79	32%
Subtotal	245	100%	245	100%	241	100%	245	100%	250	100%
Female adult or adolescent										
Injection drug use	4	8%	3	7%	1	2%	3	8%	-	-
Heterosexual contact	27	56%	25	57%	36	65%	15	38%	16	32%
Other/unknown	17	35%	16	36%	18	33%	21	54%	34	68%
Subtotal	48	100%	44	100%	55	100%	39	100%	50	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	100%	2	100%	2	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	2	100%	-	-	1	100%
Total	296		291		298		284		301	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 157. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Cleveland Part A, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	111	82%	110	83%	125	78%	137	86%	133	81%	3,334	87%
Females	25	18%	22	17%	35	22%	22	14%	32	19%	485	13%
Age at AIDS diagnosis (yr)												
<13	2	1%	2	2%	1	1%	-	-	1	1%	20	1%
13-14	1	1%	-	-	-	-	-	-	-	-	4	<1%
15-19	5	4%	1	1%	3	2%	3	2%	3	2%	32	1%
20-24	8	6%	12	9%	12	8%	16	10%	18	11%	198	5%
25-29	15	11%	13	10%	27	17%	16	10%	14	8%	510	13%
30-34	12	9%	22	17%	19	12%	24	15%	29	18%	833	22%
35-39	22	16%	22	17%	19	12%	21	13%	34	21%	760	20%
40-44	31	23%	22	17%	23	14%	32	20%	24	15%	632	17%
45-49	20	15%	19	14%	29	18%	21	13%	16	10%	413	11%
50-54	12	9%	13	10%	14	9%	10	6%	10	6%	225	6%
55-64	7	5%	5	4%	10	6%	13	8%	16	10%	166	4%
65+	1	1%	1	1%	3	2%	3	2%	-	-	25	1%
Race/Ethnicity												
White, not Hispanic	70	51%	64	48%	69	43%	83	52%	76	46%	2,471	65%
Black/African American, not Hispanic	51	38%	56	42%	80	50%	70	44%	70	42%	1,185	31%
Hispanic/Latino	9	7%	5	4%	6	4%	4	3%	10	6%	104	3%
Asian/Pacific Islander	1	1%	1	1%	1	1%	-	-	2	1%	14	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	1%	2	<1%
Unknown	5	4%	6	5%	4	3%	2	1%	6	4%	43	1%
Race/Sex												
White, not Hispanic Males	63	46%	60	45%	60	38%	81	51%	68	41%	2,310	60%
White, not Hispanic Females	7	5%	4	3%	9	6%	2	1%	8	5%	161	4%
Black/African American, not Hispanic Males	34	25%	40	30%	56	35%	51	32%	50	30%	888	23%
Black/African American, not Hispanic Females	17	13%	16	12%	24	15%	19	12%	20	12%	297	8%
Hispanic/Latino Males	8	6%	4	3%	6	4%	4	3%	7	4%	86	2%
Hispanic/Latina Females	1	1%	1	1%	-	-	-	-	3	2%	18	<1%
Asian/Pacific Islander Males	1	1%	1	1%	-	-	-	-	2	1%	12	<1%
Asian/Pacific Islander Females	-	-	-	-	1	1%	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	1	1%	2	<1%
Unknown	5	4%	6	5%	4	3%	2	1%	6	4%	43	1%
Total	136		132		160		159		165		3,819	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part A Columbus Planning Region

Table 158. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Cleveland Part A, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	74	68%	74	68%	89	71%	79	58%	81	61%	2,397	72%
Injection drug use (IDU)	1	1%	3	3%	3	2%	-	-	3	2%	183	6%
Male-to-male sexual contact and IDU	6	6%	7	6%	3	2%	6	4%	4	3%	134	4%
Heterosexual contact	16	15%	15	14%	16	13%	9	7%	5	4%	177	5%
Other/unknown	12	11%	10	9%	14	11%	43	31%	40	30%	431	13%
Subtotal	109	100%	109	100%	125	100%	137	100%	133	100%	3,322	100%
Female adult or adolescent												
Injection drug use	4	17%	1	5%	2	6%	2	9%	-	-	72	15%
Heterosexual contact	14	58%	12	60%	20	59%	11	50%	14	45%	256	54%
Other/unknown	6	25%	7	35%	12	35%	9	41%	17	55%	142	30%
Subtotal	24	100%	20	100%	34	100%	22	100%	31	100%	470	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	100%	2	67%	1	100%	-	-	1	100%	21	78%
Other/unknown	-	-	1	33%	-	-	-	-	-	-	6	22%
Subtotal	3	100%	3	100%	1	100%	-	-	1	100%	27	100%
Total	136		132		160		159		165		3,819	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 159. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Cleveland Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	341.9	3,283	77%	1,641	77%	1,642	78%
Females	92.8	961	23%	499	23%	462	22%
Age at diagnosis (yr)							
<13	4.5	14	<1%	9	<1%	5	<1%
13-14	11.2	6	<1%	4	<1%	2	<1%
15-19	18.3	25	1%	19	1%	6	<1%
20-24	128.7	152	4%	119	6%	33	2%
25-29	216.2	259	6%	192	9%	67	3%
30-34	270.0	317	7%	213	10%	104	5%
35-39	292.3	334	8%	199	9%	135	6%
40-44	486.6	641	15%	324	15%	317	15%
45-49	538.5	783	18%	349	16%	434	21%
50-54	466.5	752	18%	305	14%	447	21%
55-64	286.8	784	18%	331	15%	453	22%
65+	56.9	177	4%	76	4%	101	5%
Race/Ethnicity							
White, not Hispanic	106.4	1,499	35%	744	35%	755	36%
Black/African American, not Hispanic	554.9	2,272	54%	1,156	54%	1,116	53%
Hispanic/Latino	365.6	369	9%	170	8%	199	9%
Asian/Pacific Islander	37.0	15	<1%	11	1%	4	<1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	85	2%	57	3%	28	1%
Race/Sex							
White, not Hispanic Males	190.9	1,306	31%	637	30%	669	32%
White, not Hispanic Females	26.6	193	5%	107	5%	86	4%
Black/African American, not Hispanic Males	874.6	1,650	39%	837	39%	813	39%
Black/African American, not Hispanic Females	281.8	622	15%	319	15%	303	14%
Hispanic/Latino Males	488.6	247	6%	114	5%	133	6%
Hispanic/Latina Females	242.1	122	3%	56	3%	66	3%
Asian/Pacific Islander Males	56.8	11	<1%	8	<1%	3	<1%
Asian/Pacific Islander Females	*	4	<1%	3	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	85	2%	57	3%	28	1%
Total	212.6	4,244		2,140		2,104	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 160. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Cleveland Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,150	66%	1,034	64%	1,116	68%
Injection drug use (IDU)	189	6%	68	4%	121	7%
Male-to-male sexual contact and IDU	147	5%	44	3%	103	6%
Heterosexual contact	238	7%	114	7%	124	8%
Other/unknown	530	16%	364	22%	166	10%
Subtotal	3,254	100%	1,624	100%	1,630	100%
Female adult or adolescent						
Injection drug use	122	13%	39	8%	83	18%
Heterosexual contact	572	61%	278	58%	294	65%
Other/unknown	238	26%	166	34%	72	16%
Subtotal	932	100%	483	100%	449	100%
Child (<13 yrs at diagnosis)						
Perinatal	50	86%	27	82%	23	92%
Other/unknown	8	14%	6	18%	2	8%
Subtotal	58	100%	33	100%	25	100%
Total	4,244		2,140		2,104	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 161. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Cleveland Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	2,598	77%	2,782	77%	2,948	77%	3,102	77%	3,283	77%
Females	766	23%	825	23%	881	23%	915	23%	961	23%
Age at diagnosis (yr)										
<13	11	<1%	12	<1%	14	<1%	14	<1%	14	<1%
13-14	5	<1%	6	<1%	6	<1%	6	<1%	6	<1%
15-19	18	1%	18	<1%	18	<1%	21	1%	25	1%
20-24	30	1%	46	1%	70	2%	109	3%	152	4%
25-29	139	4%	176	5%	204	5%	230	6%	259	6%
30-34	187	6%	222	6%	258	7%	281	7%	317	7%
35-39	247	7%	276	8%	296	8%	318	8%	334	8%
40-44	524	16%	559	15%	598	16%	615	15%	641	15%
45-49	689	20%	715	20%	740	19%	759	19%	783	18%
50-54	658	20%	692	19%	710	19%	729	18%	752	18%
55-64	690	21%	715	20%	742	19%	761	19%	784	18%
65+	166	5%	170	5%	173	5%	174	4%	177	4%
Race/Ethnicity										
White, not Hispanic	1,224	36%	1,306	36%	1,376	36%	1,430	36%	1,499	35%
Black/African American, not Hispanic	1,751	52%	1,887	52%	2,017	53%	2,131	53%	2,272	54%
Hispanic/Latino	300	9%	323	9%	339	9%	355	9%	369	9%
Asian/Pacific Islander	13	<1%	13	<1%	14	<1%	15	<1%	15	<1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	72	2%	74	2%	79	2%	82	2%	85	2%
Race/Sex										
White, not Hispanic Males	1,065	32%	1,138	32%	1,199	31%	1,248	31%	1,306	31%
White, not Hispanic Females	159	5%	168	5%	177	5%	182	5%	193	5%
Black/African American, not Hispanic Males	1,264	38%	1,359	38%	1,450	38%	1,540	38%	1,650	39%
Black/African American, not Hispanic Females	487	14%	528	15%	567	15%	591	15%	622	15%
Hispanic/Latino Males	200	6%	214	6%	225	6%	236	6%	247	6%
Hispanic/Latina Females	100	3%	109	3%	114	3%	119	3%	122	3%
Asian/Pacific Islander Males	10	<1%	10	<1%	10	<1%	11	<1%	11	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	4	<1%	4	<1%	4	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	72	2%	74	2%	79	2%	82	2%	85	2%
Total	3,364		3,607		3,829		4,017		4,244	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 162. Reported persons living with a diagnosis of HIV infection by year and transmission category, Cleveland Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,681	65%	1,813	66%	1,928	66%	2,027	66%	2,150	66%
Injection drug use (IDU)	176	7%	180	7%	186	6%	186	6%	189	6%
Male-to-male sexual contact and IDU	131	5%	140	5%	144	5%	145	5%	147	5%
Heterosexual contact	195	8%	211	8%	229	8%	232	8%	238	7%
Other/unknown	388	15%	411	15%	432	15%	483	16%	530	16%
Subtotal	2,571	100%	2,755	100%	2,919	100%	3,073	100%	3,254	100%
Female adult or adolescent										
Injection drug use	108	15%	111	14%	114	13%	118	13%	122	13%
Heterosexual contact	472	64%	514	65%	547	64%	556	63%	572	61%
Other/unknown	159	22%	171	21%	191	22%	212	24%	238	26%
Subtotal	739	100%	796	100%	852	100%	886	100%	932	100%
Child (<13 yrs at diagnosis)										
Perinatal	47	87%	48	86%	50	86%	50	86%	50	86%
Other/unknown	7	13%	8	14%	8	14%	8	14%	8	14%
Subtotal	54	100%	56	100%	58	100%	58	100%	58	100%
Total	3,364		3,607		3,829		4,017		4,244	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 163. Reported diagnoses of HIV infection by year of diagnosis selected characteristics, Cleveland Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	20.5	197	83%	20.6	198	76%	18.2	175	75%	16.5	158	82%	19.2	184	80%
Females	3.9	40	17%	6.2	64	24%	5.7	59	25%	3.4	35	18%	4.4	46	20%
Age at diagnosis (yr)															
<13	*	1	<1%	*	2	1%	*	2	1%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
15-19	8.8	12	5%	8.8	12	5%	8.0	11	5%	13.9	19	10%	10.2	14	6%
20-24	34.7	41	17%	31.3	37	14%	25.4	30	13%	29.6	35	18%	34.7	41	18%
25-29	20.9	25	11%	30.9	37	14%	32.5	39	17%	21.7	26	13%	27.5	33	14%
30-34	18.7	22	9%	25.6	30	11%	15.3	18	8%	22.1	26	13%	25.6	30	13%
35-39	32.4	37	16%	29.8	34	13%	33.3	38	16%	17.5	20	10%	12.3	14	6%
40-44	26.6	35	15%	22.0	29	11%	19.7	26	11%	12.1	16	8%	22.0	29	13%
45-49	19.9	29	12%	23.4	34	13%	19.9	29	12%	15.1	22	11%	20.6	30	13%
50-54	11.2	18	8%	14.9	24	9%	12.4	20	9%	8.7	14	7%	11.8	19	8%
55-64	3.7	10	4%	7.7	21	8%	6.2	17	7%	5.1	14	7%	6.6	18	8%
65+	2.3	7	3%	*	2	1%	*	4	2%	*	-	-	*	2	1%
Race/Ethnicity															
White, not Hispanic	5.7	80	34%	6.3	89	34%	5.4	76	32%	4.1	58	30%	5.0	70	30%
Black/African American, not Hispanic	33.0	135	57%	35.7	146	56%	33.2	136	58%	28.1	115	60%	34.9	143	62%
Hispanic/Latino	19.8	20	8%	23.8	24	9%	15.9	16	7%	15.9	16	8%	13.9	14	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	1	<1%	*	1	1%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	3	1%	*	5	2%	*	3	2%	*	3	1%
Race/Sex															
White, not Hispanic Males	10.7	73	31%	11.5	79	30%	9.6	66	28%	7.6	52	27%	8.6	59	26%
White, not Hispanic Females	1.0	7	3%	1.4	10	4%	1.4	10	4%	0.8	6	3%	1.5	11	5%
Black/African American, not Hispanic Males	56.7	107	45%	54.6	103	39%	50.4	95	41%	48.2	91	47%	59.4	112	49%
Black/African American, not Hispanic Females	12.7	28	12%	19.5	43	16%	18.6	41	18%	10.9	24	12%	14.0	31	13%
Hispanic/Latino Males	29.7	15	6%	27.7	14	5%	21.8	11	5%	21.8	11	6%	21.8	11	5%
Hispanic/Latina Females	9.9	5	2%	19.8	10	4%	9.9	5	2%	9.9	5	3%	*	3	1%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	3	1%	*	5	2%	*	3	2%	*	3	1%
Total	11.9	237		13.1	262		11.7	234		9.7	193		11.5	230	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 164. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Cleveland Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	138	70%	142	72%	113	65%	102	65%	125	68%
Injection drug use (IDU)	4	2%	3	2%	7	4%	-	-	3	2%
Male-to-male sexual contact and IDU	4	2%	8	4%	5	3%	1	1%	2	1%
Heterosexual contact	21	11%	17	9%	18	10%	3	2%	6	3%
Other/unknown	29	15%	28	14%	30	17%	52	33%	48	26%
Subtotal	196	100%	198	100%	173	100%	158	100%	184	100%
Female adult or adolescent										
Injection drug use	3	8%	4	6%	3	5%	4	11%	4	9%
Heterosexual contact	28	70%	45	73%	33	56%	9	26%	16	35%
Other/unknown	9	23%	13	21%	23	39%	22	63%	26	57%
Subtotal	40	100%	62	100%	59	100%	35	100%	46	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	1	50%	2	100%	-	-	-	-
Other/unknown	-	-	1	50%	-	-	-	-	-	-
Subtotal	1	100%	2	100%	2	100%	-	-	-	-
Total	237		262		234		193		230	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 165. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Cleveland Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	80	78%	85	75%	88	73%	73	74%	76	78%	3,774	82%
Females	23	22%	28	25%	33	27%	26	26%	22	22%	825	18%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	48	1%
13-14	1	1%	-	-	-	-	-	-	-	-	3	<1%
15-19	4	4%	-	-	3	2%	3	3%	2	2%	36	1%
20-24	11	11%	9	8%	8	7%	12	12%	6	6%	218	5%
25-29	6	6%	13	12%	8	7%	11	11%	12	12%	582	13%
30-34	8	8%	15	13%	13	11%	12	12%	10	10%	904	20%
35-39	18	17%	16	14%	24	20%	12	12%	5	5%	1,002	22%
40-44	18	17%	22	19%	16	13%	13	13%	19	19%	741	16%
45-49	18	17%	17	15%	22	18%	19	19%	24	24%	522	11%
50-54	11	11%	9	8%	12	10%	11	11%	9	9%	252	5%
55-64	2	2%	11	10%	14	12%	6	6%	9	9%	231	5%
65+	6	6%	1	1%	1	1%	-	-	2	2%	59	1%
Race/Ethnicity												
White, not Hispanic	42	41%	35	31%	38	31%	29	29%	27	28%	1,841	40%
Black/African American, not Hispanic	51	50%	59	52%	69	57%	60	61%	66	67%	2,337	51%
Hispanic/Latino	10	10%	17	15%	11	9%	9	9%	4	4%	374	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	8	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	6	<1%
Unknown	-	-	2	2%	3	2%	1	1%	1	1%	33	1%
Race/Sex												
White, not Hispanic Males	35	34%	32	28%	34	28%	24	24%	23	23%	1,687	37%
White, not Hispanic Females	7	7%	3	3%	4	3%	5	5%	4	4%	154	3%
Black/African American, not Hispanic Males	38	37%	44	39%	46	38%	43	43%	48	49%	1,794	39%
Black/African American, not Hispanic Females	13	13%	15	13%	23	19%	17	17%	18	18%	543	12%
Hispanic/Latino Males	7	7%	8	7%	6	5%	5	5%	4	4%	259	6%
Hispanic/Latina Females	3	3%	9	8%	5	4%	4	4%	-	-	115	3%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	7	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	3	<1%
Unknown	-	-	2	2%	3	2%	1	1%	1	1%	33	1%
Total	103		113		121		99		98		4,599	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cleveland Consortium 1

Table 166. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Cleveland Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	54	68%	58	68%	53	60%	48	66%	44	59%	2,545	68%
Injection drug use (IDU)	3	4%	2	2%	6	7%	2	3%	1	1%	380	10%
Male-to-male sexual contact and IDU	2	3%	7	8%	6	7%	3	4%	2	3%	281	7%
Heterosexual contact	11	14%	11	13%	12	14%	3	4%	3	4%	208	6%
Other/unknown	10	13%	7	8%	11	13%	17	23%	25	33%	336	9%
Subtotal	80	100%	85	100%	88	100%	73	100%	75	100%	3,750	100%
Female adult or adolescent												
Injection drug use	6	27%	2	7%	4	12%	5	21%	1	5%	193	24%
Heterosexual contact	14	64%	22	79%	24	73%	9	38%	9	41%	466	59%
Other/unknown	2	9%	4	14%	5	15%	10	42%	12	55%	132	17%
Subtotal	22	100%	28	100%	33	100%	24	100%	22	100%	791	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	-	-	2	100%	1	100%	50	86%
Other/unknown	-	-	-	-	-	-	-	-	-	-	8	14%
Subtotal	1	100%	-	-	-	-	2	100%	1	100%	58	100%
Total	103		113		121		99		98		4,599	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 167. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Columbus Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	381.7	3,417	82%	1,914	81%	1,503	83%
Females	81.2	754	18%	445	19%	309	17%
Age at diagnosis (yr)							
<13	4.0	13	<1%	6	<1%	7	<1%
13-14	*	-	-	-	-	-	-
15-19	18.6	23	1%	17	1%	6	<1%
20-24	145.6	199	5%	153	6%	46	3%
25-29	228.2	317	8%	230	10%	87	5%
30-34	295.7	394	9%	263	11%	131	7%
35-39	355.2	440	11%	250	11%	190	10%
40-44	487.6	636	15%	350	15%	286	16%
45-49	629.9	815	20%	422	18%	393	22%
50-54	469.1	610	15%	300	13%	310	17%
55-64	291.4	606	15%	303	13%	303	17%
65+	60.9	119	3%	66	3%	53	3%
Race/Ethnicity							
White, not Hispanic	172.4	2,369	57%	1,357	58%	1,012	56%
Black/African American, not Hispanic	552.6	1,522	36%	848	36%	674	37%
Hispanic/Latino	216.8	150	4%	74	3%	76	4%
Asian/Pacific Islander	36.3	22	1%	13	1%	9	<1%
American Indian/Alaska Native	*	4	<1%	3	<1%	1	<1%
Unknown	*	105	3%	65	3%	40	2%
Race/Sex							
White, not Hispanic Males	316.8	2,139	51%	1,212	51%	927	51%
White, not Hispanic Females	32.9	230	6%	145	6%	85	5%
Black/African American, not Hispanic Males	793.1	1,049	25%	575	24%	474	26%
Black/African American, not Hispanic Females	330.4	473	11%	273	12%	200	11%
Hispanic/Latino Males	331.2	121	3%	61	3%	60	3%
Hispanic/Latina Females	88.8	29	1%	13	1%	16	1%
Asian/Pacific Islander Males	57.3	17	<1%	10	<1%	7	<1%
Asian/Pacific Islander Females	16.1	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native Males	*	3	<1%	3	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Unknown	*	105	3%	65	3%	40	2%
Total	228.8	4,172		2,360		1,812	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 168. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Columbus Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,314	68%	1,289	68%	1,025	68%
Injection drug use (IDU)	75	2%	21	1%	54	4%
Male-to-male sexual contact and IDU	93	3%	40	2%	53	4%
Heterosexual contact	219	6%	84	4%	135	9%
Other/unknown	700	21%	470	25%	230	15%
Subtotal	3,401	100%	1,904	100%	1,497	100%
Female adult or adolescent						
Injection drug use	61	8%	28	6%	33	11%
Heterosexual contact	394	54%	213	49%	181	61%
Other/unknown	280	38%	195	45%	85	28%
Subtotal	735	100%	436	100%	299	100%
Child (<13 yrs at diagnosis)						
Perinatal	29	83%	16	84%	13	81%
Other/unknown	6	17%	3	16%	3	19%
Subtotal	35	100%	19	100%	16	100%
Total	4,172		2,360		1,812	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 169. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Columbus Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	2,484	81%	2,710	82%	2,938	81%	3,173	82%	3,417	82%
Females	568	19%	613	18%	668	19%	706	18%	754	18%
Age at diagnosis (yr)										
<13	9	<1%	11	<1%	12	<1%	12	<1%	13	<1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	10	<1%	10	<1%	13	<1%	15	<1%	23	1%
20-24	37	1%	56	2%	101	3%	147	4%	199	5%
25-29	116	4%	164	5%	219	6%	267	7%	317	8%
30-34	224	7%	261	8%	304	8%	347	9%	394	9%
35-39	291	10%	328	10%	364	10%	400	10%	440	11%
40-44	491	16%	529	16%	570	16%	603	16%	636	15%
45-49	705	23%	740	22%	763	21%	791	20%	815	20%
50-54	533	17%	558	17%	573	16%	591	15%	610	15%
55-64	536	18%	561	17%	576	16%	591	15%	606	15%
65+	101	3%	106	3%	112	3%	116	3%	119	3%
Race/Ethnicity										
White, not Hispanic	1,797	59%	1,943	58%	2,088	58%	2,227	57%	2,369	57%
Black/African American, not Hispanic	1,064	35%	1,170	35%	1,289	36%	1,401	36%	1,522	36%
Hispanic/Latino	100	3%	109	3%	119	3%	132	3%	150	4%
Asian/Pacific Islander	14	<1%	16	<1%	18	<1%	20	1%	22	1%
American Indian/Alaska Native	3	<1%	4	<1%	4	<1%	4	<1%	4	<1%
Unknown	75	2%	82	2%	89	2%	96	2%	105	3%
Race/Sex										
White, not Hispanic Males	1,621	53%	1,753	53%	1,883	52%	2,010	52%	2,139	51%
White, not Hispanic Females	176	6%	190	6%	205	6%	217	6%	230	6%
Black/African American, not Hispanic Males	710	23%	787	24%	870	24%	959	25%	1,049	25%
Black/African American, not Hispanic Females	354	12%	383	12%	419	12%	442	11%	473	11%
Hispanic/Latino Males	77	3%	86	3%	95	3%	107	3%	121	3%
Hispanic/Latina Females	23	1%	23	1%	24	1%	25	1%	29	1%
Asian/Pacific Islander Males	11	<1%	13	<1%	14	<1%	15	<1%	17	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	4	<1%	5	<1%	5	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	75	2%	82	2%	89	2%	96	2%	105	3%
Total	3,053		3,324		3,607		3,880		4,172	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 170. Reported persons living with a diagnosis of HIV infection by year and transmission category, Columbus Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,656	67%	1,833	68%	2,002	69%	2,159	68%	2,314	68%
Injection drug use (IDU)	68	3%	69	3%	70	2%	70	2%	75	2%
Male-to-male sexual contact and IDU	66	3%	73	3%	80	3%	88	3%	93	3%
Heterosexual contact	161	7%	187	7%	205	7%	215	7%	219	6%
Other/unknown	517	21%	532	20%	565	19%	625	20%	700	21%
Subtotal	2,468	100%	2,694	100%	2,922	100%	3,157	100%	3,401	100%
Female adult or adolescent										
Injection drug use	54	10%	57	10%	58	9%	61	9%	61	8%
Heterosexual contact	301	54%	327	55%	363	56%	378	55%	394	54%
Other/unknown	199	36%	213	36%	229	35%	249	36%	280	38%
Subtotal	554	100%	597	100%	650	100%	688	100%	735	100%
Child (<13 yrs at diagnosis)										
Perinatal	24	80%	26	81%	28	82%	28	82%	29	83%
Other/unknown	6	20%	6	19%	6	18%	6	18%	6	17%
Subtotal	30	100%	32	100%	34	100%	34	100%	35	100%
Total	3,053		3,324		3,607		3,880		4,172	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 171. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Columbus Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	27.3	244	83%	27.4	245	84%	26.8	240	81%	27.4	245	86%	27.7	248	83%
Females	5.4	50	17%	5.0	46	16%	6.1	57	19%	4.2	39	14%	5.4	50	17%
Age at diagnosis (yr)															
<13	*	3	1%	*	2	1%	*	2	1%	*	-	-	*	1	<1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	10.5	13	4%	10.5	13	4%	21.8	27	9%	10.5	13	5%	12.9	16	5%
20-24	30.0	41	14%	35.8	49	17%	38.0	52	18%	43.9	60	21%	41.0	56	19%
25-29	41.8	58	20%	26.6	37	13%	41.0	57	19%	31.0	43	15%	33.1	46	15%
30-34	19.5	26	9%	33.0	44	15%	26.3	35	12%	30.0	40	14%	34.5	46	15%
35-39	38.7	48	16%	31.5	39	13%	33.1	41	14%	29.9	37	13%	34.7	43	14%
40-44	41.4	54	18%	32.2	42	14%	27.6	36	12%	27.6	36	13%	26.1	34	11%
45-49	17.8	23	8%	22.4	29	10%	13.9	18	6%	20.1	26	9%	16.2	21	7%
50-54	14.6	19	6%	19.2	25	9%	10.8	14	5%	9.2	12	4%	15.4	20	7%
55-64	3.8	8	3%	2.9	6	2%	5.8	12	4%	7.2	15	5%	5.8	12	4%
65+	*	1	<1%	2.6	5	2%	*	3	1%	*	2	1%	*	3	1%
Race/Ethnicity															
White, not Hispanic	11.5	158	54%	11.3	156	54%	11.1	153	52%	10.5	145	51%	10.6	146	49%
Black/African American, not Hispanic	40.3	111	38%	41.8	115	40%	45.4	125	42%	42.5	117	41%	44.7	123	41%
Hispanic/Latino	20.2	14	5%	14.5	10	3%	14.5	10	3%	18.8	13	5%	26.0	18	6%
Asian/Pacific Islander	*	3	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
American Indian/Alaska Native	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	8	3%	*	7	2%	*	7	2%	*	7	2%	*	9	3%
Race/Sex															
White, not Hispanic Males	21.2	143	49%	21.0	142	49%	20.4	138	46%	19.7	133	47%	19.5	132	44%
White, not Hispanic Females	2.1	15	5%	2.0	14	5%	2.1	15	5%	1.7	12	4%	2.0	14	5%
Black/African American, not Hispanic Males	59.7	79	27%	65.0	86	30%	65.8	87	29%	70.3	93	33%	68.8	91	31%
Black/African American, not Hispanic Females	22.4	32	11%	20.3	29	10%	26.5	38	13%	16.8	24	8%	22.4	32	11%
Hispanic/Latino Males	35.6	13	4%	24.6	9	3%	24.6	9	3%	32.8	12	4%	38.3	14	5%
Hispanic/Latina Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	4	1%
Asian/Pacific Islander Males	*	3	1%	*	2	1%	*	1	<1%	*	1	<1%	*	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	1	<1%	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
Unknown	*	8	3%	*	7	2%	*	7	2%	*	7	2%	*	9	3%
Total	16.1	294		16.0	291		16.3	297		15.6	284		16.3	298	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 172. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Columbus Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	187	77%	191	78%	177	74%	162	66%	156	63%
Injection drug use (IDU)	6	2%	2	1%	1	<1%	-	-	5	2%
Male-to-male sexual contact and IDU	8	3%	9	4%	7	3%	8	3%	5	2%
Heterosexual contact	12	5%	27	11%	18	8%	10	4%	4	2%
Other/unknown	30	12%	16	7%	37	15%	65	27%	78	31%
Subtotal	243	100%	245	100%	240	100%	245	100%	248	100%
Female adult or adolescent										
Injection drug use	4	8%	3	7%	1	2%	3	8%	-	-
Heterosexual contact	27	56%	25	57%	36	65%	15	38%	16	33%
Other/unknown	17	35%	16	36%	18	33%	21	54%	33	67%
Subtotal	48	100%	44	100%	55	100%	39	100%	49	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	100%	2	100%	2	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	2	100%	2	100%	-	-	1	100%
Total	294		291		297		284		298	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 173. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Columbus Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	110	81%	110	83%	125	78%	137	86%	132	81%	3,319	87%
Females	25	19%	22	17%	35	22%	22	14%	31	19%	482	13%
Age at AIDS diagnosis (yr)												
<13	2	1%	2	2%	1	1%	-	-	1	1%	20	1%
13-14	1	1%	-	-	-	-	-	-	-	-	4	<1%
15-19	5	4%	1	1%	3	2%	3	2%	3	2%	32	1%
20-24	8	6%	12	9%	12	8%	16	10%	18	11%	198	5%
25-29	15	11%	13	10%	27	17%	16	10%	14	9%	508	13%
30-34	12	9%	22	17%	19	12%	24	15%	29	18%	831	22%
35-39	21	16%	22	17%	19	12%	21	13%	33	20%	756	20%
40-44	31	23%	22	17%	23	14%	32	20%	23	14%	627	16%
45-49	20	15%	19	14%	29	18%	21	13%	16	10%	411	11%
50-54	12	9%	13	10%	14	9%	10	6%	10	6%	223	6%
55-64	7	5%	5	4%	10	6%	13	8%	16	10%	165	4%
65+	1	1%	1	1%	3	2%	3	2%	-	-	25	1%
Race/Ethnicity												
White, not Hispanic	69	51%	64	48%	69	43%	83	52%	74	45%	2,453	65%
Black/African American, not Hispanic	51	38%	56	42%	80	50%	70	44%	70	43%	1,185	31%
Hispanic/Latino	9	7%	5	4%	6	4%	4	3%	10	6%	104	3%
Asian/Pacific Islander	1	1%	1	1%	1	1%	-	-	2	1%	14	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	1%	2	<1%
Unknown	5	4%	6	5%	4	3%	2	1%	6	4%	43	1%
Race/Sex												
White, not Hispanic Males	62	46%	60	45%	60	38%	81	51%	67	41%	2,295	60%
White, not Hispanic Females	7	5%	4	3%	9	6%	2	1%	7	4%	158	4%
Black/African American, not Hispanic Males	34	25%	40	30%	56	35%	51	32%	50	31%	888	23%
Black/African American, not Hispanic Females	17	13%	16	12%	24	15%	19	12%	20	12%	297	8%
Hispanic/Latino Males	8	6%	4	3%	6	4%	4	3%	7	4%	86	2%
Hispanic/Latina Females	1	1%	1	1%	-	-	-	-	3	2%	18	<1%
Asian/Pacific Islander Males	1	1%	1	1%	-	-	-	-	2	1%	12	<1%
Asian/Pacific Islander Females	-	-	-	-	1	1%	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	1	1%	2	<1%
Unknown	5	4%	6	5%	4	3%	2	1%	6	4%	43	1%
Total	135		132		160		159		163		3,801	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Columbus Consortium 2

Table 174. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Columbus Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	74	69%	74	68%	89	71%	79	58%	80	61%	2,386	72%
Injection drug use (IDU)	1	1%	3	3%	3	2%	-	-	3	2%	182	6%
Male-to-male sexual contact and IDU	6	6%	7	6%	3	2%	6	4%	4	3%	134	4%
Heterosexual contact	16	15%	15	14%	16	13%	9	7%	5	4%	175	5%
Other/unknown	11	10%	10	9%	14	11%	43	31%	40	30%	430	13%
Subtotal	108	100%	109	100%	125	100%	137	100%	132	100%	3,307	100%
Female adult or adolescent												
Injection drug use	4	17%	1	5%	2	6%	2	9%	-	-	71	15%
Heterosexual contact	14	58%	12	60%	20	59%	11	50%	14	47%	255	55%
Other/unknown	6	25%	7	35%	12	35%	9	41%	16	53%	141	30%
Subtotal	24	100%	20	100%	34	100%	22	100%	30	100%	467	100%
Child (<13 yrs at diagnosis)												
Perinatal	3	100%	2	67%	1	100%	-	-	1	100%	21	78%
Other/unknown	-	-	1	33%	-	-	-	-	-	-	6	22%
Subtotal	3	100%	3	100%	1	100%	-	-	1	100%	27	100%
Total	135		132		160		159		163		3,801	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 175. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Cincinnati Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	262.8	2,233	79%	1,066	77%	1,167	81%
Females	66.6	595	21%	327	23%	268	19%
Age at diagnosis (yr)							
<13	*	3	<1%	2	<1%	1	<1%
13-14	*	1	<1%	-	-	1	<1%
15-19	23.9	29	1%	25	2%	4	<1%
20-24	123.1	146	5%	106	8%	40	3%
25-29	189.5	214	8%	141	10%	73	5%
30-34	189.6	209	7%	116	8%	93	6%
35-39	288.2	304	11%	169	12%	135	9%
40-44	309.2	366	13%	168	12%	198	14%
45-49	428.1	546	19%	237	17%	309	22%
50-54	332.8	445	16%	212	15%	233	16%
55-64	205.8	449	16%	164	12%	285	20%
65+	52.2	116	4%	53	4%	63	4%
Race/Ethnicity							
White, not Hispanic	92.2	1,275	45%	632	45%	643	45%
Black/African American, not Hispanic	591.2	1,442	51%	703	50%	739	51%
Hispanic/Latino	128.6	60	2%	29	2%	31	2%
Asian/Pacific Islander	34.4	13	<1%	9	1%	4	<1%
American Indian/Alaska Native	*	2	<1%	-	-	2	<1%
Unknown	*	36	1%	20	1%	16	1%
Race/Sex							
White, not Hispanic Males	159.8	1,082	38%	511	37%	571	40%
White, not Hispanic Females	27.4	193	7%	121	9%	72	5%
Black/African American, not Hispanic Males	928.4	1,060	37%	509	37%	551	38%
Black/African American, not Hispanic Females	294.5	382	14%	194	14%	188	13%
Hispanic/Latino Males	199.9	49	2%	22	2%	27	2%
Hispanic/Latina Females	49.7	11	<1%	7	1%	4	<1%
Asian/Pacific Islander Males	60.7	11	<1%	8	1%	3	<1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	*	2	<1%	-	-	2	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	36	1%	20	1%	16	1%
Total	162.3	2,828		1,393		1,435	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 176. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Cincinnati Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,484	67%	666	63%	818	70%
Injection drug use (IDU)	72	3%	32	3%	40	3%
Male-to-male sexual contact and IDU	85	4%	31	3%	54	5%
Heterosexual contact	142	6%	59	6%	83	7%
Other/unknown	440	20%	272	26%	168	14%
Subtotal	2,223	100%	1,060	100%	1,163	100%
Female adult or adolescent						
Injection drug use	52	9%	23	7%	29	11%
Heterosexual contact	314	54%	160	50%	154	59%
Other/unknown	219	37%	139	43%	80	30%
Subtotal	585	100%	322	100%	263	100%
Child (<13 yrs at diagnosis)						
Perinatal	15	75%	9	82%	6	67%
Other/unknown	5	25%	2	18%	3	33%
Subtotal	20	100%	11	100%	9	100%
Total	2,828		1,393		1,435	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 177. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Cincinnati Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	1,711	80%	1,824	80%	1,961	80%	2,097	79%	2,233	79%
Females	422	20%	458	20%	504	20%	546	21%	595	21%
Age at diagnosis (yr)										
<13	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	8	<1%	8	<1%	9	<1%	18	1%	29	1%
20-24	22	1%	39	2%	76	3%	110	4%	146	5%
25-29	85	4%	120	5%	153	6%	186	7%	214	8%
30-34	127	6%	141	6%	166	7%	188	7%	209	7%
35-39	223	10%	238	10%	259	11%	279	11%	304	11%
40-44	301	14%	317	14%	329	13%	348	13%	366	13%
45-49	471	22%	495	22%	512	21%	532	20%	546	19%
50-54	379	18%	394	17%	410	17%	425	16%	445	16%
55-64	403	19%	416	18%	436	18%	441	17%	449	16%
65+	110	5%	110	5%	111	5%	112	4%	116	4%
Race/Ethnicity										
White, not Hispanic	1,040	49%	1,093	48%	1,147	47%	1,207	46%	1,275	45%
Black/African American, not Hispanic	1,018	48%	1,103	48%	1,222	50%	1,333	50%	1,442	51%
Hispanic/Latino	38	2%	48	2%	53	2%	55	2%	60	2%
Asian/Pacific Islander	9	<1%	9	<1%	10	<1%	11	<1%	13	<1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	26	1%	27	1%	31	1%	35	1%	36	1%
Race/Sex										
White, not Hispanic Males	892	42%	937	41%	980	40%	1,026	39%	1,082	38%
White, not Hispanic Females	148	7%	156	7%	167	7%	181	7%	193	7%
Black/African American, not Hispanic Males	757	35%	815	36%	900	37%	985	37%	1,060	37%
Black/African American, not Hispanic Females	261	12%	288	13%	322	13%	348	13%	382	14%
Hispanic/Latino Males	30	1%	39	2%	43	2%	45	2%	49	2%
Hispanic/Latina Females	8	<1%	9	<1%	10	<1%	10	<1%	11	<1%
Asian/Pacific Islander Males	8	<1%	8	<1%	9	<1%	10	<1%	11	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Males	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	26	1%	27	1%	31	1%	35	1%	36	1%
Total	2,133		2,282		2,465		2,643		2,828	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 178. Reported persons living with a diagnosis of HIV infection by year and transmission category, Cincinnati Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,140	67%	1,215	67%	1,313	67%	1,406	67%	1,484	67%
Injection drug use (IDU)	66	4%	69	4%	71	4%	71	3%	72	3%
Male-to-male sexual contact and IDU	72	4%	76	4%	79	4%	82	4%	85	4%
Heterosexual contact	108	6%	119	7%	128	7%	138	7%	142	6%
Other/unknown	315	19%	335	18%	360	18%	390	19%	440	20%
Subtotal	1,701	100%	1,814	100%	1,951	100%	2,087	100%	2,223	100%
Female adult or adolescent										
Injection drug use	35	8%	39	9%	45	9%	49	9%	52	9%
Heterosexual contact	254	62%	273	61%	289	59%	303	57%	314	54%
Other/unknown	123	30%	136	30%	160	32%	184	34%	219	37%
Subtotal	412	100%	448	100%	494	100%	536	100%	585	100%
Child (<13 yrs at diagnosis)										
Perinatal	15	75%	15	75%	15	75%	15	75%	15	75%
Other/unknown	5	25%	5	25%	5	25%	5	25%	5	25%
Subtotal	20	100%	20	100%	20	100%	20	100%	20	100%
Total	2,133		2,282		2,465		2,643		2,828	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 179. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Cincinnati Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	13.8	117	79%	14.6	124	76%	17.1	145	74%	16.2	138	75%	16.4	139	74%
Females	3.6	32	21%	4.4	39	24%	5.6	50	26%	5.0	45	25%	5.6	50	26%
Age at diagnosis (yr)															
<13	*	1	1%	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	8.2	10	7%	7.4	9	6%	14.0	17	9%	15.7	19	10%	9.9	12	6%
20-24	20.2	24	16%	33.7	40	25%	37.1	44	23%	32.9	39	21%	35.4	42	22%
25-29	18.6	21	14%	14.2	16	10%	25.7	29	15%	28.3	32	17%	23.9	27	14%
30-34	18.1	20	13%	12.7	14	9%	22.7	25	13%	16.3	18	10%	25.4	28	15%
35-39	18.0	19	13%	19.0	20	12%	13.3	14	7%	19.0	20	11%	13.3	14	7%
40-44	15.2	18	12%	25.3	30	18%	11.8	14	7%	16.9	20	11%	16.9	20	11%
45-49	11.8	15	10%	11.0	14	9%	14.9	19	10%	18.0	23	13%	15.7	20	11%
50-54	7.5	10	7%	9.7	13	8%	13.5	18	9%	6.7	9	5%	12.0	16	8%
55-64	4.1	9	6%	3.2	7	4%	6.4	14	7%	*	1	1%	3.2	7	4%
65+	*	2	1%	*	-	-	*	1	1%	*	2	1%	*	3	2%
Race/Ethnicity															
White, not Hispanic	4.2	58	39%	4.1	57	35%	4.2	58	30%	4.5	62	34%	5.1	71	38%
Black/African American, not Hispanic	33.6	82	55%	38.5	94	58%	51.7	126	65%	46.3	113	62%	45.1	110	58%
Hispanic/Latino	10.7	5	3%	23.6	11	7%	12.9	6	3%	*	2	1%	10.7	5	3%
Asian/Pacific Islander	*	2	1%	*	-	-	*	1	1%	*	1	1%	*	2	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	1	1%	*	4	2%	*	5	3%	*	1	1%
Race/Sex															
White, not Hispanic Males	6.6	45	30%	7.1	48	29%	6.8	46	24%	6.9	47	26%	8.7	59	31%
White, not Hispanic Females	1.8	13	9%	1.3	9	6%	1.7	12	6%	2.1	15	8%	1.7	12	6%
Black/African American, not Hispanic Males	57.8	66	44%	56.9	65	40%	78.0	89	46%	75.3	86	47%	65.7	75	40%
Black/African American, not Hispanic Females	12.3	16	11%	22.4	29	18%	28.5	37	19%	20.8	27	15%	27.0	35	19%
Hispanic/Latino Males	*	3	2%	40.8	10	6%	20.4	5	3%	*	2	1%	*	4	2%
Hispanic/Latina Females	*	2	1%	*	1	1%	*	1	1%	*	-	-	*	1	1%
Asian/Pacific Islander Males	*	2	1%	*	-	-	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	1%	*	1	1%	*	4	2%	*	5	3%	*	1	1%
Total	8.5	149		9.4	163		11.2	195		10.5	183		10.8	189	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 180. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Cincinnati Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	84	72%	79	64%	102	70%	95	69%	79	57%
Injection drug use (IDU)	2	2%	3	2%	2	1%	-	-	1	1%
Male-to-male sexual contact and IDU	2	2%	4	3%	3	2%	3	2%	4	3%
Heterosexual contact	20	17%	11	9%	9	6%	10	7%	4	3%
Other/unknown	9	8%	27	22%	29	20%	30	22%	51	37%
Subtotal	117	100%	124	100%	145	100%	138	100%	139	100%
Female adult or adolescent										
Injection drug use	1	3%	4	10%	7	14%	4	9%	3	6%
Heterosexual contact	25	81%	20	51%	17	34%	14	31%	11	22%
Other/unknown	5	16%	15	38%	26	52%	27	60%	36	72%
Subtotal	31	100%	39	100%	50	100%	45	100%	50	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-
Total	149		163		195		183		189	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 181. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Cincinnati Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	54	81%	86	77%	93	73%	85	80%	77	74%	2,438	86%
Females	13	19%	26	23%	35	27%	21	20%	27	26%	383	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	18	1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	4	4%	6	5%	1	1%	2	2%	27	1%
20-24	6	9%	9	8%	13	10%	9	8%	21	20%	145	5%
25-29	7	10%	10	9%	20	16%	25	24%	10	10%	411	15%
30-34	10	15%	8	7%	11	9%	15	14%	19	18%	580	21%
35-39	13	19%	21	19%	18	14%	11	10%	11	11%	589	21%
40-44	11	16%	23	21%	26	20%	12	11%	7	7%	454	16%
45-49	10	15%	16	14%	15	12%	14	13%	10	10%	264	9%
50-54	3	4%	13	12%	13	10%	9	8%	12	12%	176	6%
55-64	6	9%	8	7%	5	4%	7	7%	10	10%	120	4%
65+	1	1%	-	-	1	1%	3	3%	2	2%	36	1%
Race/Ethnicity												
White, not Hispanic	27	40%	44	39%	41	32%	38	36%	40	38%	1,500	53%
Black/African American, not Hispanic	38	57%	62	55%	77	60%	66	62%	60	58%	1,253	44%
Hispanic/Latino	2	3%	4	4%	5	4%	1	1%	2	2%	41	1%
Asian/Pacific Islander	-	-	-	-	1	1%	-	-	-	-	7	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Unknown	-	-	2	2%	4	3%	1	1%	2	2%	18	1%
Race/Sex												
White, not Hispanic Males	23	34%	39	35%	36	28%	33	31%	31	30%	1,390	49%
White, not Hispanic Females	4	6%	5	4%	5	4%	5	5%	9	9%	110	4%
Black/African American, not Hispanic Males	30	45%	41	37%	51	40%	50	47%	43	41%	991	35%
Black/African American, not Hispanic Females	8	12%	21	19%	26	20%	16	15%	17	16%	262	9%
Hispanic/Latino Males	1	1%	4	4%	3	2%	1	1%	2	2%	35	1%
Hispanic/Latina Females	1	1%	-	-	2	2%	-	-	-	-	6	<1%
Asian/Pacific Islander Males	-	-	-	-	1	1%	-	-	-	-	6	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	2	2%	4	3%	1	1%	2	2%	18	1%
Total	67		112		128		106		104		2,821	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Cincinnati Consortium 3

Table 182. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Cincinnati Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	39	72%	57	66%	59	63%	61	72%	49	64%	1,748	72%
Injection drug use (IDU)	1	2%	1	1%	2	2%	-	-	-	-	105	4%
Male-to-male sexual contact and IDU	1	2%	1	1%	3	3%	4	5%	2	3%	138	6%
Heterosexual contact	10	19%	12	14%	12	13%	3	4%	2	3%	112	5%
Other/unknown	3	6%	15	17%	17	18%	17	20%	24	31%	320	13%
Subtotal	54	100%	86	100%	93	100%	85	100%	77	100%	2,423	100%
Female adult or adolescent												
Injection drug use	-	-	1	4%	5	14%	1	5%	3	12%	56	15%
Heterosexual contact	10	77%	20	77%	18	51%	5	24%	8	31%	203	54%
Other/unknown	3	23%	5	19%	12	34%	15	71%	15	58%	116	31%
Subtotal	13	100%	26	100%	35	100%	21	100%	26	100%	375	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	1	100%	17	74%
Other/unknown	-	-	-	-	-	-	-	-	-	-	6	26%
Subtotal	-	-	-	-	-	-	-	-	1	100%	23	100%
Total	67		112		128		106		104		2,821	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 183. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Dayton Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	239.8	1,204	79%	577	77%	627	80%
Females	61.4	328	21%	176	23%	152	20%
Age at diagnosis (yr)							
<13	3.6	6	<1%	5	1%	1	<1%
13-14	*	3	<1%	3	<1%	-	-
15-19	13.8	10	1%	7	1%	3	<1%
20-24	115.4	83	5%	67	9%	16	2%
25-29	172.0	110	7%	71	9%	39	5%
30-34	213.2	130	8%	76	10%	54	7%
35-39	262.3	150	10%	84	11%	66	8%
40-44	294.0	192	13%	82	11%	110	14%
45-49	395.3	284	19%	111	15%	173	22%
50-54	311.7	242	16%	117	16%	125	16%
55-64	186.8	258	17%	103	14%	155	20%
65+	39.5	64	4%	27	4%	37	5%
Race/Ethnicity							
White, not Hispanic	91.7	765	50%	346	46%	419	54%
Black/African American, not Hispanic	511.7	708	46%	375	50%	333	43%
Hispanic/Latino	91.7	21	1%	9	1%	12	2%
Asian/Pacific Islander	46.9	8	1%	4	1%	4	1%
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	28	2%	18	2%	10	1%
Race/Sex							
White, not Hispanic Males	160.7	653	43%	289	38%	364	47%
White, not Hispanic Females	26.2	112	7%	57	8%	55	7%
Black/African American, not Hispanic Males	778.9	505	33%	264	35%	241	31%
Black/African American, not Hispanic Females	276.1	203	13%	111	15%	92	12%
Hispanic/Latino Males	159.4	19	1%	9	1%	10	1%
Hispanic/Latina Females	*	2	<1%	-	-	2	<1%
Asian/Pacific Islander Males	88.4	7	<1%	4	1%	3	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-
Unknown	*	28	2%	18	2%	10	1%
Total	147.9	1,532		753		779	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 184. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Dayton Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	863	72%	389	68%	474	76%
Injection drug use (IDU)	28	2%	15	3%	13	2%
Male-to-male sexual contact and IDU	40	3%	14	2%	26	4%
Heterosexual contact	76	6%	29	5%	47	8%
Other/unknown	187	16%	125	22%	62	10%
Subtotal	1,194	100%	572	100%	622	100%
Female adult or adolescent						
Injection drug use	35	11%	21	12%	14	9%
Heterosexual contact	200	62%	94	55%	106	71%
Other/unknown	86	27%	56	33%	30	20%
Subtotal	321	100%	171	100%	150	100%
Child (<13 yrs at diagnosis)						
Perinatal	13	76%	9	90%	4	57%
Other/unknown	4	24%	1	10%	3	43%
Subtotal	17	100%	10	100%	7	100%
Total	1,532		753		779	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 185. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Dayton Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	932	78%	989	78%	1,042	78%	1,121	79%	1,204	79%
Females	260	22%	275	22%	290	22%	305	21%	328	21%
Age at diagnosis (yr)										
<13	2	<1%	3	<1%	3	<1%	6	<1%	6	<1%
13-14	1	<1%	1	<1%	2	<1%	3	<1%	3	<1%
15-19	3	<1%	3	<1%	3	<1%	7	<1%	10	1%
20-24	11	1%	18	1%	30	2%	53	4%	83	5%
25-29	60	5%	68	5%	83	6%	100	7%	110	7%
30-34	81	7%	92	7%	101	8%	116	8%	130	8%
35-39	120	10%	128	10%	135	10%	137	10%	150	10%
40-44	158	13%	168	13%	170	13%	181	13%	192	13%
45-49	252	21%	264	21%	269	20%	275	19%	284	19%
50-54	206	17%	213	17%	223	17%	230	16%	242	16%
55-64	239	20%	247	20%	252	19%	254	18%	258	17%
65+	59	5%	59	5%	61	5%	64	4%	64	4%
Race/Ethnicity										
White, not Hispanic	636	53%	667	53%	690	52%	723	51%	765	50%
Black/African American, not Hispanic	511	43%	548	43%	592	44%	650	46%	708	46%
Hispanic/Latino	17	1%	19	2%	19	1%	19	1%	21	1%
Asian/Pacific Islander	6	1%	7	1%	7	1%	7	<1%	8	1%
American Indian/Alaska Native	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	20	2%	21	2%	22	2%	25	2%	28	2%
Race/Sex										
White, not Hispanic Males	541	45%	567	45%	587	44%	619	43%	653	43%
White, not Hispanic Females	95	8%	100	8%	103	8%	104	7%	112	7%
Black/African American, not Hispanic Males	357	30%	385	30%	417	31%	462	32%	505	33%
Black/African American, not Hispanic Females	154	13%	163	13%	175	13%	188	13%	203	13%
Hispanic/Latino Males	15	1%	17	1%	17	1%	17	1%	19	1%
Hispanic/Latina Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Asian/Pacific Islander Males	5	<1%	6	<1%	6	<1%	6	<1%	7	<1%
Asian/Pacific Islander Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	20	2%	21	2%	22	2%	25	2%	28	2%
Total	1,192		1,264		1,332		1,426		1,532	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 186. Reported persons living with a diagnosis of HIV infection by year and transmission category, Dayton Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	676	73%	720	73%	754	73%	817	74%	863	72%
Injection drug use (IDU)	24	3%	25	3%	27	3%	27	2%	28	2%
Male-to-male sexual contact and IDU	38	4%	38	4%	39	4%	40	4%	40	3%
Heterosexual contact	59	6%	61	6%	64	6%	67	6%	76	6%
Other/unknown	130	14%	139	14%	151	15%	160	14%	187	16%
Subtotal	927	100%	983	100%	1,035	100%	1,111	100%	1,194	100%
Female adult or adolescent										
Injection drug use	30	12%	31	12%	32	11%	32	11%	35	11%
Heterosexual contact	170	67%	179	67%	186	65%	191	64%	200	62%
Other/unknown	54	21%	59	22%	66	23%	75	25%	86	27%
Subtotal	254	100%	269	100%	284	100%	298	100%	321	100%
Child (<13 yrs at diagnosis)										
Perinatal	9	82%	9	75%	10	77%	13	76%	13	76%
Other/unknown	2	18%	3	25%	3	23%	4	24%	4	24%
Subtotal	11	100%	12	100%	13	100%	17	100%	17	100%
Total	1,192		1,264		1,332		1,426		1,532	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 187. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Dayton Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	12.7	64	74%	12.2	61	80%	11.6	58	77%	16.3	82	85%	16.9	85	79%
Females	4.1	22	26%	2.8	15	20%	3.2	17	23%	2.8	15	15%	4.3	23	21%
Age at diagnosis (yr)															
<13	*	1	1%	*	1	1%	*	1	1%	*	4	4%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	2%	8.3	6	8%	9.7	7	9%	19.4	14	14%	9.7	7	6%
20-24	19.5	14	16%	11.1	8	11%	22.2	16	21%	32.0	23	24%	40.3	29	27%
25-29	14.1	9	10%	17.2	11	14%	15.6	10	13%	25.0	16	16%	17.2	11	10%
30-34	19.7	12	14%	16.4	10	13%	14.8	9	12%	13.1	8	8%	23.0	14	13%
35-39	19.2	11	13%	15.7	9	12%	*	3	4%	10.5	6	6%	21.0	12	11%
40-44	21.4	14	16%	18.4	12	16%	7.7	5	7%	13.8	9	9%	15.3	10	9%
45-49	12.5	9	10%	9.7	7	9%	13.9	10	13%	8.4	6	6%	15.3	11	10%
50-54	11.6	9	10%	9.0	7	9%	7.7	6	8%	7.7	6	6%	12.9	10	9%
55-64	*	2	2%	*	4	5%	5.8	8	11%	*	2	2%	*	3	3%
65+	*	3	3%	*	1	1%	*	-	-	*	3	3%	*	1	1%
Race/Ethnicity															
White, not Hispanic	4.1	34	40%	4.1	34	45%	3.1	26	35%	4.2	35	36%	5.0	42	39%
Black/African American, not Hispanic	34.0	47	55%	27.5	38	50%	34.7	48	64%	42.6	59	61%	43.4	60	56%
Hispanic/Latino	*	2	2%	*	2	3%	*	-	-	*	-	-	*	2	2%
Asian/Pacific Islander	*	1	1%	*	1	1%	*	-	-	*	-	-	*	1	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	2%	*	1	1%	*	1	1%	*	3	3%	*	3	3%
Race/Sex															
White, not Hispanic Males	7.4	30	35%	7.1	29	38%	5.7	23	31%	8.4	34	35%	8.4	34	31%
White, not Hispanic Females	*	4	5%	1.2	5	7%	*	3	4%	*	1	1%	1.9	8	7%
Black/African American, not Hispanic Males	47.8	31	36%	44.7	29	38%	52.4	34	45%	70.9	46	47%	69.4	45	42%
Black/African American, not Hispanic Females	21.8	16	19%	12.2	9	12%	19.0	14	19%	17.7	13	13%	20.4	15	14%
Hispanic/Latino Males	*	2	2%	*	2	3%	*	-	-	*	-	-	*	2	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	-	-	*	-	-	*	1	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	2%	*	1	1%	*	1	1%	*	3	3%	*	3	3%
Total	8.3	86		7.3	76		7.2	75		9.4	97		10.4	108	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 188. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Dayton Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	46	72%	47	78%	36	63%	64	81%	46	54%
Injection drug use (IDU)	4	6%	1	2%	2	4%	1	1%	1	1%
Male-to-male sexual contact and IDU	-	-	-	-	1	2%	1	1%	-	-
Heterosexual contact	7	11%	2	3%	3	5%	3	4%	9	11%
Other/unknown	7	11%	10	17%	15	26%	10	13%	29	34%
Subtotal	64	100%	60	100%	57	100%	79	100%	85	100%
Female adult or adolescent										
Injection drug use	2	10%	1	7%	1	6%	-	-	3	13%
Heterosexual contact	14	67%	9	60%	7	41%	5	36%	9	39%
Other/unknown	5	24%	5	33%	9	53%	9	64%	11	48%
Subtotal	21	100%	15	100%	17	100%	14	100%	23	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	1	100%	3	75%	-	-
Other/unknown	-	-	1	100%	-	-	1	25%	-	-
Subtotal	1	100%	1	100%	1	100%	4	100%	-	-
Total	86		76		75		97		108	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 189. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Dayton Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	47	78%	44	80%	34	79%	32	74%	33	85%	1,369	85%
Females	13	22%	11	20%	9	21%	11	26%	6	15%	246	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	1	2%	-	-	16	1%
13-14	-	-	-	-	-	-	-	-	-	-	3	<1%
15-19	-	-	1	2%	1	2%	3	7%	2	5%	22	1%
20-24	6	10%	7	13%	5	12%	3	7%	5	13%	88	5%
25-29	7	12%	6	11%	7	16%	4	9%	5	13%	201	12%
30-34	5	8%	6	11%	5	12%	8	19%	4	10%	303	19%
35-39	7	12%	7	13%	3	7%	2	5%	3	8%	315	20%
40-44	16	27%	11	20%	7	16%	7	16%	6	15%	295	18%
45-49	6	10%	5	9%	4	9%	8	19%	7	18%	181	11%
50-54	6	10%	7	13%	6	14%	4	9%	4	10%	99	6%
55-64	2	3%	5	9%	5	12%	2	5%	3	8%	66	4%
65+	5	8%	-	-	-	-	1	2%	-	-	26	2%
Race/Ethnicity												
White, not Hispanic	28	47%	25	45%	17	40%	18	42%	20	51%	994	62%
Black/African American, not Hispanic	30	50%	28	51%	25	58%	24	56%	16	41%	589	36%
Hispanic/Latino	-	-	-	-	-	-	-	-	1	3%	16	1%
Asian/Pacific Islander	-	-	2	4%	-	-	-	-	-	-	5	<1%
American Indian/Alaska Native	-	-	-	-	0	<1%	-	-	-	-	1	<1%
Unknown	2	3%	-	-	1	2%	1	2%	2	5%	10	1%
Race/Sex												
White, not Hispanic Males	24	40%	23	42%	14	33%	16	37%	18	46%	895	55%
White, not Hispanic Females	4	7%	2	4%	3	7%	2	5%	2	5%	99	6%
Black/African American, not Hispanic Males	22	37%	19	35%	19	44%	15	35%	13	33%	446	28%
Black/African American, not Hispanic Females	8	13%	9	16%	6	14%	9	21%	3	8%	143	9%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	1	3%	15	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Males	-	-	2	4%	-	-	-	-	-	-	4	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	2	3%	-	-	1	2%	1	2%	2	5%	10	1%
Total	60		55		43		43		39		1,615	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Dayton Consortium 4

Table 190. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Dayton Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	32	68%	37	84%	25	74%	26	84%	21	64%	981	72%
Injection drug use (IDU)	1	2%	1	2%	1	3%	1	3%	-	-	59	4%
Male-to-male sexual contact and IDU	2	4%	-	-	-	-	1	3%	1	3%	87	6%
Heterosexual contact	5	11%	4	9%	2	6%	1	3%	5	15%	82	6%
Other/unknown	7	15%	2	5%	6	18%	2	6%	6	18%	145	11%
Subtotal	47	100%	44	100%	34	100%	31	100%	33	100%	1,354	100%
Female adult or adolescent												
Injection drug use	1	8%	-	-	-	-	-	-	-	-	41	17%
Heterosexual contact	11	85%	10	91%	6	67%	6	55%	4	67%	139	58%
Other/unknown	1	8%	1	9%	3	33%	5	45%	2	33%	58	24%
Subtotal	13	100%	11	100%	9	100%	11	100%	6	100%	238	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	15	65%
Other/unknown	-	-	-	-	-	-	1	100%	-	-	8	35%
Subtotal	-	-	-	-	-	-	1	100%	-	-	23	100%
Total	60		55		43		43		39		1,615	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 191. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Toledo Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	184.5	734	72%	357	71%	377	73%
Females	67.8	283	28%	147	29%	136	27%
Age at diagnosis (yr)							
<13	5.2	7	1%	3	1%	4	1%
13-14	*	1	<1%	1	<1%	-	-
15-19	13.4	8	1%	6	1%	2	<1%
20-24	53.1	33	3%	28	6%	5	1%
25-29	96.8	49	5%	35	7%	14	3%
30-34	154.2	74	7%	46	9%	28	5%
35-39	266.1	126	12%	66	13%	60	12%
40-44	359.7	182	18%	96	19%	86	17%
45-49	342.4	191	19%	77	15%	114	22%
50-54	272.4	170	17%	70	14%	100	19%
55-64	133.5	144	14%	64	13%	80	16%
65+	27.9	32	3%	12	2%	20	4%
Race/Ethnicity							
White, not Hispanic	80.5	523	51%	257	51%	266	52%
Black/African American, not Hispanic	457.8	409	40%	207	41%	202	39%
Hispanic/Latino	113.4	57	6%	23	5%	34	7%
Asian/Pacific Islander	70.4	7	1%	4	1%	3	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	21	2%	13	3%	8	2%
Race/Sex							
White, not Hispanic Males	133.2	423	42%	202	40%	221	43%
White, not Hispanic Females	30.1	100	10%	55	11%	45	9%
Black/African American, not Hispanic Males	586.3	249	24%	126	25%	123	24%
Black/African American, not Hispanic Females	341.4	160	16%	81	16%	79	15%
Hispanic/Latino Males	155.0	39	4%	15	3%	24	5%
Hispanic/Latina Females	71.8	18	2%	8	2%	10	2%
Asian/Pacific Islander Males	126.5	6	1%	4	1%	2	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	21	2%	13	3%	8	2%
Total	124.7	1,017		504		513	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 192. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Toledo Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	527	73%	252	72%	275	74%
Injection drug use (IDU)	32	4%	13	4%	19	5%
Male-to-male sexual contact and IDU	46	6%	21	6%	25	7%
Heterosexual contact	72	10%	36	10%	36	10%
Other/unknown	49	7%	30	9%	19	5%
Subtotal	726	100%	352	100%	374	100%
Female adult or adolescent						
Injection drug use	24	9%	10	7%	14	11%
Heterosexual contact	209	76%	107	75%	102	77%
Other/unknown	41	15%	25	18%	16	12%
Subtotal	274	100%	142	100%	132	100%
Child (<13 yrs at diagnosis)						
Perinatal	17	100%	10	100%	7	100%
Other/unknown	-	-	-	-	-	-
Subtotal	17	100%	10	100%	7	100%
Total	1,017		504		513	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 193. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Toledo Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	582	71%	627	71%	661	71%	690	72%	734	72%
Females	242	29%	259	29%	267	29%	273	28%	283	28%
Age at diagnosis (yr)										
<13	6	1%	7	1%	7	1%	7	1%	7	1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	5	1%	5	1%	5	1%	5	1%	8	1%
20-24	7	1%	12	1%	17	2%	24	2%	33	3%
25-29	24	3%	37	4%	41	4%	44	5%	49	5%
30-34	53	6%	60	7%	65	7%	68	7%	74	7%
35-39	97	12%	107	12%	109	12%	117	12%	126	12%
40-44	147	18%	162	18%	169	18%	174	18%	182	18%
45-49	179	22%	181	20%	185	20%	187	19%	191	19%
50-54	150	18%	154	17%	160	17%	165	17%	170	17%
55-64	124	15%	129	15%	138	15%	139	14%	144	14%
65+	31	4%	31	3%	31	3%	32	3%	32	3%
Race/Ethnicity										
White, not Hispanic	439	53%	462	52%	480	52%	500	52%	523	51%
Black/African American, not Hispanic	318	39%	350	40%	371	40%	384	40%	409	40%
Hispanic/Latino	47	6%	52	6%	54	6%	55	6%	57	6%
Asian/Pacific Islander	5	1%	7	1%	7	1%	7	1%	7	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	15	2%	15	2%	16	2%	17	2%	21	2%
Race/Sex										
White, not Hispanic Males	350	42%	370	42%	385	41%	402	42%	423	42%
White, not Hispanic Females	89	11%	92	10%	95	10%	98	10%	100	10%
Black/African American, not Hispanic Males	184	22%	204	23%	221	24%	231	24%	249	24%
Black/African American, not Hispanic Females	134	16%	146	16%	150	16%	153	16%	160	16%
Hispanic/Latino Males	31	4%	35	4%	36	4%	37	4%	39	4%
Hispanic/Latina Females	16	2%	17	2%	18	2%	18	2%	18	2%
Asian/Pacific Islander Males	5	1%	6	1%	6	1%	6	1%	6	1%
Asian/Pacific Islander Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	15	2%	15	2%	16	2%	17	2%	21	2%
Total	824		886		928		963		1,017	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 194. Reported persons living with a diagnosis of HIV infection by year and transmission category, Toledo Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	413	72%	446	72%	474	73%	496	73%	527	73%
Injection drug use (IDU)	30	5%	31	5%	31	5%	31	5%	32	4%
Male-to-male sexual contact and IDU	41	7%	43	7%	44	7%	45	7%	46	6%
Heterosexual contact	53	9%	61	10%	65	10%	68	10%	72	10%
Other/unknown	37	6%	38	6%	39	6%	42	6%	49	7%
Subtotal	574	100%	619	100%	653	100%	682	100%	726	100%
Female adult or adolescent										
Injection drug use	23	10%	24	10%	24	9%	24	9%	24	9%
Heterosexual contact	178	76%	192	77%	198	77%	203	77%	209	76%
Other/unknown	33	14%	34	14%	36	14%	37	14%	41	15%
Subtotal	234	100%	250	100%	258	100%	264	100%	274	100%
Child (<13 yrs at diagnosis)										
Perinatal	16	100%	17	100%	17	100%	17	100%	17	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	16	100%	17	100%	17	100%	17	100%	17	100%
Total	824		886		928		963		1,017	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 195. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Toledo Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	6.3	25	58%	11.3	45	73%	8.5	34	79%	8.0	32	82%	11.3	45	80%
Females	4.3	18	42%	4.1	17	27%	2.2	9	21%	1.7	7	18%	2.6	11	20%
Age at diagnosis (yr)															
<13	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	5%	*	2	3%	*	1	2%	*	2	5%	*	4	7%
20-24	8.0	5	12%	20.9	13	21%	11.3	7	16%	12.9	8	21%	16.1	10	18%
25-29	9.9	5	12%	19.7	10	16%	*	4	9%	*	1	3%	9.9	5	9%
30-34	20.8	10	23%	18.7	9	15%	*	3	7%	12.5	6	15%	16.7	8	14%
35-39	14.8	7	16%	31.7	15	24%	14.8	7	16%	21.1	10	26%	21.1	10	18%
40-44	15.8	8	19%	*	3	5%	*	4	9%	*	2	5%	9.9	5	9%
45-49	*	4	9%	*	4	6%	10.8	6	14%	*	3	8%	*	4	7%
50-54	*	2	5%	*	3	5%	8.0	5	12%	*	3	8%	9.6	6	11%
55-64	*	-	-	*	2	3%	5.6	6	14%	*	4	10%	*	4	7%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity															
White, not Hispanic	2.8	18	42%	3.5	23	37%	2.8	18	42%	3.5	23	59%	3.5	23	41%
Black/African American, not Hispanic	24.6	22	51%	35.8	32	52%	24.6	22	51%	15.7	14	36%	30.2	27	48%
Hispanic/Latino	*	2	5%	9.9	5	8%	*	2	5%	*	1	3%	*	2	4%
Asian/Pacific Islander	*	-	-	*	2	3%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	2%	*	-	-	*	1	2%	*	1	3%	*	4	7%
Race/Sex															
White, not Hispanic Males	4.4	14	33%	6.3	20	32%	4.7	15	35%	6.3	20	51%	6.6	21	38%
White, not Hispanic Females	*	4	9%	*	3	5%	*	3	7%	*	3	8%	*	2	4%
Black/African American, not Hispanic Males	21.2	9	21%	47.1	20	32%	40.0	17	40%	23.5	10	26%	44.7	19	34%
Black/African American, not Hispanic Females	27.7	13	30%	25.6	12	19%	10.7	5	12%	*	4	10%	17.1	8	14%
Hispanic/Latino Males	*	1	2%	*	4	6%	*	1	2%	*	1	3%	*	2	4%
Hispanic/Latina Females	*	1	2%	*	1	2%	*	1	2%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	2%	*	-	-	*	1	2%	*	1	3%	*	4	7%
Total	5.3	43		7.6	62		5.3	43		4.8	39		6.9	56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 196. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Toledo Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	21	84%	33	73%	28	82%	24	75%	32	71%
Injection drug use (IDU)	-	-	1	2%	-	-	-	-	1	2%
Male-to-male sexual contact and IDU	-	-	2	4%	1	3%	1	3%	1	2%
Heterosexual contact	3	12%	8	18%	4	12%	3	9%	4	9%
Other/unknown	1	4%	1	2%	1	3%	4	13%	7	16%
Subtotal	25	100%	45	100%	34	100%	32	100%	45	100%
Female adult or adolescent										
Injection drug use	-	-	1	6%	-	-	-	-	-	-
Heterosexual contact	17	94%	14	88%	6	67%	6	86%	7	64%
Other/unknown	1	6%	1	6%	3	33%	1	14%	4	36%
Subtotal	18	100%	16	100%	9	100%	7	100%	11	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	-	-	-	-	-	-
Total	43		62		43		39		56	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 197. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Toledo Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	12	48%	23	64%	27	75%	22	79%	25	83%	854	80%
Females	13	52%	13	36%	9	25%	6	21%	5	17%	214	20%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	15	1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	2	8%	1	3%	-	-	-	-	1	3%	12	1%
20-24	2	8%	2	6%	1	3%	2	7%	1	3%	63	6%
25-29	2	8%	5	14%	3	8%	-	-	3	10%	137	13%
30-34	5	20%	6	17%	5	14%	8	29%	1	3%	220	21%
35-39	4	16%	9	25%	7	19%	5	18%	5	17%	215	20%
40-44	6	24%	5	14%	10	28%	4	14%	4	13%	160	15%
45-49	3	12%	5	14%	3	8%	5	18%	6	20%	119	11%
50-54	1	4%	2	6%	3	8%	3	11%	5	17%	69	6%
55-64	-	-	1	3%	4	11%	1	4%	4	13%	46	4%
65+	-	-	-	-	-	-	-	-	-	-	11	1%
Race/Ethnicity												
White, not Hispanic	10	40%	16	44%	13	36%	18	64%	16	53%	640	60%
Black/African American, not Hispanic	13	52%	18	50%	22	61%	7	25%	12	40%	366	34%
Hispanic/Latino	2	8%	1	3%	1	3%	3	11%	2	7%	49	5%
Asian/Pacific Islander	-	-	1	3%	-	-	-	-	-	-	4	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	-	-	8	1%
Race/Sex												
White, not Hispanic Males	6	24%	14	39%	10	28%	14	50%	15	50%	559	52%
White, not Hispanic Females	4	16%	2	6%	3	8%	4	14%	1	3%	81	8%
Black/African American, not Hispanic Males	5	20%	9	25%	16	44%	5	18%	8	27%	248	23%
Black/African American, not Hispanic Females	8	32%	9	25%	6	17%	2	7%	4	13%	118	11%
Hispanic/Latino Males	1	4%	-	-	1	3%	3	11%	2	7%	36	3%
Hispanic/Latina Females	1	4%	1	3%	-	-	-	-	-	-	13	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Females	-	-	1	3%	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	-	-	8	1%
Total	25		36		36		28		30		1,068	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Toledo Consortium 5

Table 198. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Toledo Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	9	75%	18	78%	21	78%	18	82%	18	72%	615	73%
Injection drug use (IDU)	-	-	1	4%	-	-	1	5%	1	4%	48	6%
Male-to-male sexual contact and IDU	-	-	1	4%	1	4%	-	-	1	4%	64	8%
Heterosexual contact	3	25%	2	9%	4	15%	1	5%	2	8%	55	7%
Other/unknown	-	-	1	4%	1	4%	2	9%	3	12%	62	7%
Subtotal	12	100%	23	100%	27	100%	22	100%	25	100%	844	100%
Female adult or adolescent												
Injection drug use	-	-	1	8%	-	-	-	-	-	-	31	15%
Heterosexual contact	13	100%	10	83%	6	67%	5	83%	4	80%	147	71%
Other/unknown	-	-	1	8%	3	33%	1	17%	1	20%	28	14%
Subtotal	13	100%	12	100%	9	100%	6	100%	5	100%	206	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	-	-	-	-	-	-	15	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	17%
Subtotal	-	-	1	100%	-	-	-	-	-	-	18	100%
Total	25		36		36		28		30		1,068	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 199. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Akron Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	160.3	682	81%	351	79%	331	83%
Females	35.6	160	19%	91	21%	69	17%
Age at diagnosis (yr)							
<13	3.7	5	1%	5	1%	-	-
13-14	*	-	-	-	-	-	-
15-19	11.0	7	1%	6	1%	1	<1%
20-24	41.4	25	3%	20	5%	5	1%
25-29	93.8	48	6%	38	9%	10	3%
30-34	139.3	70	8%	52	12%	18	5%
35-39	148.3	75	9%	43	10%	32	8%
40-44	188.1	113	13%	59	13%	54	14%
45-49	238.6	156	19%	75	17%	81	20%
50-54	216.8	151	18%	68	15%	83	21%
55-64	130.9	156	19%	63	14%	93	23%
65+	29.0	36	4%	13	3%	23	6%
Race/Ethnicity							
White, not Hispanic	65.1	482	57%	245	55%	237	59%
Black/African American, not Hispanic	368.4	320	38%	176	40%	144	36%
Hispanic/Latino	153.2	22	3%	11	2%	11	3%
Asian/Pacific Islander	36.2	6	1%	4	1%	2	1%
American Indian/Alaska Native	*	2	<1%	2	<1%	-	-
Unknown	*	10	1%	4	1%	6	2%
Race/Sex							
White, not Hispanic Males	116.3	421	50%	214	48%	207	52%
White, not Hispanic Females	16.1	61	7%	31	7%	30	8%
Black/African American, not Hispanic Males	569.2	230	27%	121	27%	109	27%
Black/African American, not Hispanic Females	193.7	90	11%	55	12%	35	9%
Hispanic/Latino Males	250.0	18	2%	9	2%	9	2%
Hispanic/Latina Females	*	4	<1%	2	<1%	2	1%
Asian/Pacific Islander Males	*	3	<1%	3	1%	-	-
Asian/Pacific Islander Females	*	3	<1%	1	<1%	2	1%
American Indian/Alaska Native Males	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	1	<1%	-	-
Unknown	*	10	1%	4	1%	6	2%
Total	96.3	842		442		400	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 200. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Akron Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	447	66%	211	61%	236	72%
Injection drug use (IDU)	24	4%	7	2%	17	5%
Male-to-male sexual contact and IDU	24	4%	7	2%	17	5%
Heterosexual contact	46	7%	19	6%	27	8%
Other/unknown	133	20%	100	29%	33	10%
Subtotal	674	100%	344	100%	330	100%
Female adult or adolescent						
Injection drug use	15	9%	8	9%	7	10%
Heterosexual contact	93	59%	48	54%	45	65%
Other/unknown	50	32%	33	37%	17	25%
Subtotal	158	100%	89	100%	69	100%
Child (<13 yrs at diagnosis)						
Perinatal	10	100%	9	100%	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	10	100%	9	100%	1	100%
Total	842		442		400	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 201. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Akron Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	538	81%	571	81%	610	81%	636	81%	682	81%
Females	127	19%	134	19%	142	19%	151	19%	160	19%
Age at diagnosis (yr)										
<13	3	<1%	3	<1%	3	<1%	3	<1%	5	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	4	1%	4	1%	5	1%	5	1%	7	1%
20-24	5	1%	8	1%	12	2%	15	2%	25	3%
25-29	13	2%	22	3%	37	5%	39	5%	48	6%
30-34	45	7%	53	8%	58	8%	67	9%	70	8%
35-39	57	9%	62	9%	66	9%	69	9%	75	9%
40-44	94	14%	97	14%	100	13%	104	13%	113	13%
45-49	132	20%	138	20%	148	20%	152	19%	156	19%
50-54	139	21%	140	20%	144	19%	147	19%	151	18%
55-64	142	21%	146	21%	147	20%	151	19%	156	19%
65+	31	5%	32	5%	32	4%	35	4%	36	4%
Race/Ethnicity										
White, not Hispanic	382	57%	402	57%	430	57%	451	57%	482	57%
Black/African American, not Hispanic	252	38%	270	38%	288	38%	299	38%	320	38%
Hispanic/Latino	18	3%	20	3%	21	3%	21	3%	22	3%
Asian/Pacific Islander	5	1%	5	1%	5	1%	6	1%	6	1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
Unknown	7	1%	7	1%	7	1%	8	1%	10	1%
Race/Sex										
White, not Hispanic Males	339	51%	357	51%	379	50%	395	50%	421	50%
White, not Hispanic Females	43	6%	45	6%	51	7%	56	7%	61	7%
Black/African American, not Hispanic Males	175	26%	188	27%	204	27%	213	27%	230	27%
Black/African American, not Hispanic Females	77	12%	82	12%	84	11%	86	11%	90	11%
Hispanic/Latino Males	14	2%	16	2%	17	2%	17	2%	18	2%
Hispanic/Latina Females	4	1%	4	1%	4	1%	4	1%	4	<1%
Asian/Pacific Islander Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
Asian/Pacific Islander Females	2	<1%	2	<1%	2	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	1	<1%	1	<1%
Unknown	7	1%	7	1%	7	1%	8	1%	10	1%
Total	665		705		752		787		842	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 202. Reported persons living with a diagnosis of HIV infection by year and transmission category, Akron Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	356	67%	386	68%	408	68%	423	67%	447	66%
Injection drug use (IDU)	20	4%	20	4%	22	4%	23	4%	24	4%
Male-to-male sexual contact and IDU	22	4%	22	4%	22	4%	24	4%	24	4%
Heterosexual contact	39	7%	41	7%	44	7%	44	7%	46	7%
Other/unknown	95	18%	96	17%	108	18%	116	18%	133	20%
Subtotal	532	100%	565	100%	604	100%	630	100%	674	100%
Female adult or adolescent										
Injection drug use	12	10%	14	11%	15	11%	15	10%	15	9%
Heterosexual contact	80	64%	81	61%	87	62%	91	61%	93	59%
Other/unknown	33	26%	37	28%	38	27%	43	29%	50	32%
Subtotal	125	100%	132	100%	140	100%	149	100%	158	100%
Child (<13 yrs at diagnosis)										
Perinatal	8	100%	8	100%	8	100%	8	100%	10	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	8	100%	8	100%	8	100%	8	100%	10	100%
Total	665		705		752		787		842	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 203. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Akron Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	9.4	40	73%	8.2	35	83%	9.6	41	82%	6.3	27	75%	11.3	48	84%
Females	3.3	15	27%	1.6	7	17%	2.0	9	18%	2.0	9	25%	2.0	9	16%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	2	4%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	3	5%	*	3	7%	*	2	4%	*	-	-	*	3	5%
20-24	*	4	7%	14.9	9	21%	24.8	15	30%	8.3	5	14%	21.5	13	23%
25-29	15.6	8	15%	15.6	8	19%	13.7	7	14%	*	2	6%	15.6	8	14%
30-34	*	4	7%	*	4	10%	*	4	8%	17.9	9	25%	*	3	5%
35-39	15.8	8	15%	*	3	7%	*	2	4%	*	4	11%	15.8	8	14%
40-44	23.3	14	25%	*	4	10%	13.3	8	16%	*	2	6%	10.0	6	11%
45-49	9.2	6	11%	7.6	5	12%	12.2	8	16%	7.6	5	14%	*	3	5%
50-54	7.2	5	9%	*	2	5%	*	1	2%	7.2	5	14%	8.6	6	11%
55-64	*	3	5%	*	4	10%	*	3	6%	*	1	3%	4.2	5	9%
65+	*	-	-	*	-	-	*	-	-	*	3	8%	*	-	-
Race/Ethnicity															
White, not Hispanic	2.6	19	35%	3.0	22	52%	3.9	29	58%	2.8	21	58%	4.3	32	56%
Black/African American, not Hispanic	36.8	32	58%	20.7	18	43%	23.0	20	40%	12.7	11	31%	25.3	22	39%
Hispanic/Latino	*	3	5%	*	2	5%	*	1	2%	*	-	-	*	1	2%
Asian/Pacific Islander	*	1	2%	*	-	-	*	-	-	*	2	6%	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	1	3%	*	2	4%
Race/Sex															
White, not Hispanic Males	4.4	16	29%	5.5	20	48%	6.4	23	46%	4.4	16	44%	7.5	27	47%
White, not Hispanic Females	*	3	5%	*	2	5%	1.6	6	12%	1.3	5	14%	1.3	5	9%
Black/African American, not Hispanic Males	52.0	21	38%	32.2	13	31%	42.1	17	34%	22.3	9	25%	44.5	18	32%
Black/African American, not Hispanic Females	23.7	11	20%	10.8	5	12%	*	3	6%	*	2	6%	*	4	7%
Hispanic/Latino Males	*	3	5%	*	2	5%	*	1	2%	*	-	-	*	1	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-
Asian/Pacific Islander Females	*	1	2%	*	-	-	*	-	-	*	1	3%	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	1	3%	*	2	4%
Total	6.3	55		4.8	42		5.7	50		4.1	36		6.5	57	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 204. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Akron Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	28	70%	32	91%	22	54%	15	56%	24	52%
Injection drug use (IDU)	2	5%	-	-	2	5%	1	4%	1	2%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	2	7%	-	-
Heterosexual contact	5	13%	2	6%	4	10%	1	4%	2	4%
Other/unknown	5	13%	1	3%	13	32%	8	30%	19	41%
Subtotal	40	100%	35	100%	41	100%	27	100%	46	100%
Female adult or adolescent										
Injection drug use	1	7%	2	29%	1	11%	-	-	-	-
Heterosexual contact	11	73%	1	14%	7	78%	4	44%	2	22%
Other/unknown	3	20%	4	57%	1	11%	5	56%	7	78%
Subtotal	15	100%	7	100%	9	100%	9	100%	9	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	2	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	2	100%
Total	55		42		50		36		57	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 205. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Akron Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	21	81%	14	78%	20	87%	11	73%	14	82%	742	87%
Females	5	19%	4	22%	3	13%	4	27%	3	18%	115	13%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	1	<1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	1	4%	-	-	-	-	1	7%	-	-	5	1%
20-24	-	-	2	11%	2	9%	1	7%	3	18%	43	5%
25-29	2	8%	1	6%	3	13%	1	7%	3	18%	105	12%
30-34	2	8%	2	11%	5	22%	3	20%	3	18%	187	22%
35-39	5	19%	2	11%	1	4%	-	-	1	6%	154	18%
40-44	7	27%	2	11%	5	22%	1	7%	3	18%	162	19%
45-49	2	8%	5	28%	3	13%	2	13%	2	12%	90	11%
50-54	5	19%	4	22%	2	9%	4	27%	2	12%	62	7%
55-64	2	8%	-	-	2	9%	1	7%	-	-	41	5%
65+	-	-	-	-	-	-	1	7%	-	-	6	1%
Race/Ethnicity												
White, not Hispanic	14	54%	6	33%	10	43%	10	67%	10	59%	543	63%
Black/African American, not Hispanic	9	35%	12	67%	10	43%	3	20%	6	35%	291	34%
Hispanic/Latino	3	12%	-	-	1	4%	-	-	-	-	14	2%
Asian/Pacific Islander	-	-	-	-	1	4%	1	7%	-	-	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	4%	1	7%	1	6%	6	1%
Race/Sex												
White, not Hispanic Males	13	50%	5	28%	9	39%	7	47%	9	53%	491	57%
White, not Hispanic Females	1	4%	1	6%	1	4%	3	20%	1	6%	52	6%
Black/African American, not Hispanic Males	5	19%	9	50%	9	39%	2	13%	4	24%	233	27%
Black/African American, not Hispanic Females	4	15%	3	17%	1	4%	1	7%	2	12%	58	7%
Hispanic/Latino Males	3	12%	-	-	1	4%	-	-	-	-	11	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	3	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	1	7%	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	1	4%	-	-	-	-	2	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	4%	1	7%	1	6%	6	1%
Total	26		18		23		15		17		857	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Akron Consortium 6

Table 206. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Akron Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	13	62%	6	43%	14	70%	7	64%	10	71%	533	72%
Injection drug use (IDU)	3	14%	-	-	3	15%	1	9%	1	7%	45	6%
Male-to-male sexual contact and IDU	-	-	1	7%	-	-	-	-	-	-	43	6%
Heterosexual contact	2	10%	3	21%	2	10%	1	9%	2	14%	39	5%
Other/unknown	3	14%	4	29%	1	5%	2	18%	1	7%	81	11%
Subtotal	21	100%	14	100%	20	100%	11	100%	14	100%	741	100%
Female adult or adolescent												
Injection drug use	1	20%	2	50%	-	-	-	-	-	-	17	15%
Heterosexual contact	3	60%	1	25%	2	67%	1	25%	1	33%	66	57%
Other/unknown	1	20%	1	25%	1	33%	3	75%	2	67%	32	28%
Subtotal	5	100%	4	100%	3	100%	4	100%	3	100%	115	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	1	100%
Total	26		18		23		15		17		857	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 207. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Youngstown Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	131.3	399	69%	207	68%	192	69%
Females	57.4	183	31%	97	32%	86	31%
Age at diagnosis (yr)							
<13	*	3	1%	3	1%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	2	<1%	2	1%	-	-
20-24	68.6	25	4%	17	6%	8	3%
25-29	121.3	40	7%	29	10%	11	4%
30-34	93.6	32	5%	21	7%	11	4%
35-39	179.0	63	11%	39	13%	24	9%
40-44	181.4	71	12%	36	12%	35	13%
45-49	223.8	97	17%	43	14%	54	19%
50-54	228.5	116	20%	53	17%	63	23%
55-64	115.2	107	18%	47	15%	60	22%
65+	23.7	26	4%	14	5%	12	4%
Race/Ethnicity							
White, not Hispanic	49.4	263	45%	130	43%	133	48%
Black/African American, not Hispanic	418.9	251	43%	127	42%	124	45%
Hispanic/Latino	230.0	38	7%	23	8%	15	5%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	2	<1%	1	<1%	1	<1%
Unknown	*	28	5%	23	8%	5	2%
Race/Sex							
White, not Hispanic Males	77.7	201	35%	97	32%	104	37%
White, not Hispanic Females	22.7	62	11%	33	11%	29	10%
Black/African American, not Hispanic Males	533.2	156	27%	82	27%	74	27%
Black/African American, not Hispanic Females	309.8	95	16%	45	15%	50	18%
Hispanic/Latino Males	233.7	21	4%	11	4%	10	4%
Hispanic/Latina Females	225.6	17	3%	12	4%	5	2%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	28	5%	23	8%	5	2%
Total	93.4	582		304		278	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 208. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Youngstown Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	206	52%	94	46%	112	59%
Injection drug use (IDU)	24	6%	7	3%	17	9%
Male-to-male sexual contact and IDU	11	3%	4	2%	7	4%
Heterosexual contact	49	12%	23	11%	26	14%
Other/unknown	103	26%	75	37%	28	15%
Subtotal	393	100%	203	100%	190	100%
Female adult or adolescent						
Injection drug use	18	10%	5	5%	13	15%
Heterosexual contact	98	56%	45	49%	53	63%
Other/unknown	60	34%	42	46%	18	21%
Subtotal	176	100%	92	100%	84	100%
Child (<13 yrs at diagnosis)						
Perinatal	11	85%	8	89%	3	75%
Other/unknown	2	15%	1	11%	1	25%
Subtotal	13	100%	9	100%	4	100%
Total	582		304		278	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 209. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Youngstown Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	310	67%	328	67%	348	67%	373	68%	399	69%
Females	150	33%	161	33%	170	33%	173	32%	183	31%
Age at diagnosis (yr)										
<13	3	1%	3	1%	3	1%	3	1%	3	1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
20-24	5	1%	6	1%	8	2%	15	3%	25	4%
25-29	18	4%	22	4%	30	6%	33	6%	40	7%
30-34	20	4%	24	5%	29	6%	30	5%	32	5%
35-39	49	11%	53	11%	59	11%	60	11%	63	11%
40-44	58	13%	62	13%	64	12%	68	12%	71	12%
45-49	83	18%	85	17%	88	17%	93	17%	97	17%
50-54	99	22%	108	22%	110	21%	114	21%	116	20%
55-64	98	21%	99	20%	100	19%	103	19%	107	18%
65+	25	5%	25	5%	25	5%	25	5%	26	4%
Race/Ethnicity										
White, not Hispanic	210	46%	223	46%	238	46%	249	46%	263	45%
Black/African American, not Hispanic	196	43%	210	43%	221	43%	235	43%	251	43%
Hispanic/Latino	32	7%	33	7%	35	7%	36	7%	38	7%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
Unknown	21	5%	22	4%	23	4%	25	5%	28	5%
Race/Sex										
White, not Hispanic Males	162	35%	171	35%	182	35%	192	35%	201	35%
White, not Hispanic Females	48	10%	52	11%	56	11%	57	10%	62	11%
Black/African American, not Hispanic Males	116	25%	124	25%	131	25%	143	26%	156	27%
Black/African American, not Hispanic Females	80	17%	86	18%	90	17%	92	17%	95	16%
Hispanic/Latino Males	18	4%	18	4%	19	4%	20	4%	21	4%
Hispanic/Latina Females	14	3%	15	3%	16	3%	16	3%	17	3%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	21	5%	22	4%	23	4%	25	5%	28	5%
Total	460		489		518		546		582	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 210. Reported persons living with a diagnosis of HIV infection by year and transmission category, Youngstown Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	159	52%	168	52%	181	53%	195	53%	206	52%
Injection drug use (IDU)	20	7%	22	7%	22	6%	24	7%	24	6%
Male-to-male sexual contact and IDU	10	3%	11	3%	11	3%	11	3%	11	3%
Heterosexual contact	37	12%	39	12%	44	13%	47	13%	49	12%
Other/unknown	78	26%	82	25%	84	25%	90	25%	103	26%
Subtotal	304	100%	322	100%	342	100%	367	100%	393	100%
Female adult or adolescent										
Injection drug use	16	11%	16	10%	16	10%	17	10%	18	10%
Heterosexual contact	79	55%	86	56%	94	58%	95	57%	98	56%
Other/unknown	48	34%	52	34%	53	33%	54	33%	60	34%
Subtotal	143	100%	154	100%	163	100%	166	100%	176	100%
Child (<13 yrs at diagnosis)										
Perinatal	11	85%	11	85%	11	85%	11	85%	11	85%
Other/unknown	2	15%	2	15%	2	15%	2	15%	2	15%
Subtotal	13	100%	13	100%	13	100%	13	100%	13	100%
Total	460		489		518		546		582	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 211. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Youngstown Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	7.2	22	73%	5.6	17	59%	7.9	24	71%	8.6	26	90%	9.2	28	74%
Females	2.5	8	27%	3.8	12	41%	3.1	10	29%	*	3	10%	3.1	10	26%
Age at diagnosis (yr)															
<13	*	1	3%	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	3%	*	-	-	*	1	3%	*	4	14%	*	4	11%
20-24	*	1	3%	*	3	10%	19.2	7	21%	*	4	14%	16.5	6	16%
25-29	*	4	13%	15.2	5	17%	*	4	12%	*	3	10%	24.3	8	21%
30-34	*	2	7%	*	2	7%	17.5	6	18%	*	-	-	*	2	5%
35-39	17.0	6	20%	14.2	5	17%	14.2	5	15%	*	3	10%	*	3	8%
40-44	12.8	5	17%	*	2	7%	*	3	9%	15.3	6	21%	*	4	11%
45-49	13.8	6	20%	18.5	8	28%	*	2	6%	*	4	14%	*	3	8%
50-54	*	2	7%	*	3	10%	*	3	9%	*	3	10%	*	3	8%
55-64	*	2	7%	*	1	3%	*	3	9%	*	2	7%	*	4	11%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
Race/Ethnicity															
White, not Hispanic	2.8	15	50%	2.4	13	45%	3.6	19	56%	2.3	12	41%	2.8	15	39%
Black/African American, not Hispanic	20.0	12	40%	21.7	13	45%	20.0	12	35%	23.4	14	48%	28.4	17	45%
Hispanic/Latino	*	1	3%	*	2	7%	*	2	6%	*	1	3%	*	2	5%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
Unknown	*	2	7%	*	1	3%	*	1	3%	*	2	7%	*	3	8%
Race/Sex															
White, not Hispanic Males	4.6	12	40%	3.5	9	31%	5.8	15	44%	4.2	11	38%	3.9	10	26%
White, not Hispanic Females	*	3	10%	*	4	14%	*	4	12%	*	1	3%	1.8	5	13%
Black/African American, not Hispanic Males	27.3	8	27%	20.5	6	21%	23.9	7	21%	41.0	12	41%	47.8	14	37%
Black/African American, not Hispanic Females	*	4	13%	22.8	7	24%	16.3	5	15%	*	2	7%	*	3	8%
Hispanic/Latino Males	*	-	-	*	1	3%	*	1	3%	*	1	3%	*	1	3%
Hispanic/Latina Females	*	1	3%	*	1	3%	*	1	3%	*	-	-	*	1	3%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	2	7%	*	1	3%	*	1	3%	*	2	7%	*	3	8%
Total	4.8	30		4.7	29		5.5	34		4.7	29		6.1	38	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 212. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Youngstown Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	15	71%	10	59%	15	63%	14	54%	11	39%
Injection drug use (IDU)	-	-	2	12%	-	-	2	8%	-	-
Male-to-male sexual contact and IDU	1	5%	-	-	-	-	-	-	-	-
Heterosexual contact	4	19%	2	12%	5	21%	3	12%	2	7%
Other/unknown	1	5%	3	18%	4	17%	7	27%	15	54%
Subtotal	21	100%	17	100%	24	100%	26	100%	28	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	1	33%	1	10%
Heterosexual contact	6	75%	7	58%	8	80%	1	33%	3	30%
Other/unknown	2	25%	5	42%	2	20%	1	33%	6	60%
Subtotal	8	100%	12	100%	10	100%	3	100%	10	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-
Total	30		29		34		29		38	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 213. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Youngstown Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	18	67%	15	75%	11	65%	12	71%	19	79%	484	77%
Females	9	33%	5	25%	6	35%	5	29%	5	21%	147	23%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	1	<1%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	2	12%	-	-	13	2%
20-24	-	-	-	-	-	-	1	6%	4	17%	23	4%
25-29	2	7%	4	20%	1	6%	1	6%	3	13%	69	11%
30-34	5	19%	2	10%	3	18%	-	-	1	4%	113	18%
35-39	4	15%	3	15%	1	6%	1	6%	2	8%	135	21%
40-44	6	22%	2	10%	4	24%	4	24%	4	17%	111	18%
45-49	5	19%	5	25%	4	24%	4	24%	2	8%	80	13%
50-54	-	-	2	10%	2	12%	3	18%	2	8%	39	6%
55-64	5	19%	1	5%	2	12%	1	6%	6	25%	39	6%
65+	-	-	1	5%	-	-	-	-	-	-	7	1%
Race/Ethnicity												
White, not Hispanic	11	41%	11	55%	13	76%	6	35%	11	46%	357	57%
Black/African American, not Hispanic	14	52%	6	30%	4	24%	9	53%	10	42%	238	38%
Hispanic/Latino	1	4%	1	5%	-	-	1	6%	1	4%	29	5%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	1	5%	-	-	-	-	-	-	1	<1%
Unknown	1	4%	1	5%	-	-	1	6%	2	8%	6	1%
Race/Sex												
White, not Hispanic Males	10	37%	8	40%	10	59%	5	29%	8	33%	305	48%
White, not Hispanic Females	1	4%	3	15%	3	18%	1	6%	3	13%	52	8%
Black/African American, not Hispanic Males	7	26%	4	20%	1	6%	5	29%	10	42%	155	25%
Black/African American, not Hispanic Females	7	26%	2	10%	3	18%	4	24%	-	-	83	13%
Hispanic/Latino Males	-	-	1	5%	-	-	1	6%	-	-	19	3%
Hispanic/Latina Females	1	4%	-	-	-	-	-	-	1	4%	10	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	1	5%	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	4%	1	5%	-	-	1	6%	2	8%	6	1%
Total	27		20		17		17		24		631	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Youngstown Consortium 7

Table 214. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Youngstown Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	11	61%	9	60%	7	64%	6	50%	8	42%	275	57%
Injection drug use (IDU)	-	-	2	13%	-	-	2	17%	-	-	45	9%
Male-to-male sexual contact and IDU	2	11%	-	-	-	-	-	-	-	-	33	7%
Heterosexual contact	3	17%	2	13%	2	18%	1	8%	4	21%	44	9%
Other/unknown	2	11%	2	13%	2	18%	3	25%	7	37%	83	17%
Subtotal	18	100%	15	100%	11	100%	12	100%	19	100%	480	100%
Female adult or adolescent												
Injection drug use	-	-	2	40%	1	17%	1	20%	2	40%	33	23%
Heterosexual contact	7	78%	3	60%	2	33%	4	80%	1	20%	71	49%
Other/unknown	2	22%	-	-	3	50%	-	-	2	40%	41	28%
Subtotal	9	100%	5	100%	6	100%	5	100%	5	100%	145	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	67%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	33%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	27		20		17		17		24		631	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 215. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Canton Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	107.5	352	81%	192	81%	160	80%
Females	24.8	85	19%	46	19%	39	20%
Age at diagnosis (yr)							
<13	*	-	-	-	-	-	-
13-14	*	1	<1%	-	-	1	1%
15-19	*	3	1%	2	1%	1	1%
20-24	63.9	26	6%	24	10%	2	1%
25-29	85.5	32	7%	25	11%	7	4%
30-34	69.4	26	6%	18	8%	8	4%
35-39	80.6	30	7%	11	5%	19	10%
40-44	194.3	82	19%	46	19%	36	18%
45-49	167.1	78	18%	35	15%	43	22%
50-54	127.5	66	15%	31	13%	35	18%
55-64	79.5	73	17%	36	15%	37	19%
65+	18.7	20	5%	10	4%	10	5%
Race/Ethnicity							
White, not Hispanic	49.3	301	69%	171	72%	130	65%
Black/African American, not Hispanic	351.0	111	25%	55	23%	56	28%
Hispanic/Latino	112.7	12	3%	4	2%	8	4%
Asian/Pacific Islander	*	2	<1%	2	1%	-	-
American Indian/Alaska Native	*	2	<1%	1	<1%	1	1%
Unknown	*	9	2%	5	2%	4	2%
Race/Sex							
White, not Hispanic Males	83.9	250	57%	141	59%	109	55%
White, not Hispanic Females	16.3	51	12%	30	13%	21	11%
Black/African American, not Hispanic Males	515.5	81	19%	41	17%	40	20%
Black/African American, not Hispanic Females	188.6	30	7%	14	6%	16	8%
Hispanic/Latino Males	175.0	10	2%	4	2%	6	3%
Hispanic/Latina Females	*	2	<1%	-	-	2	1%
Asian/Pacific Islander Males	*	1	<1%	1	<1%	-	-
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Males	*	2	<1%	1	<1%	1	1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	9	2%	5	2%	4	2%
Total	65.3	437		238		199	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 216. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Canton Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	247	70%	130	68%	117	73%
Injection drug use (IDU)	7	2%	5	3%	2	1%
Male-to-male sexual contact and IDU	12	3%	5	3%	7	4%
Heterosexual contact	21	6%	7	4%	14	9%
Other/unknown	65	18%	45	23%	20	13%
Subtotal	352	100%	192	100%	160	100%
Female adult or adolescent						
Injection drug use	8	10%	4	9%	4	11%
Heterosexual contact	48	60%	23	53%	25	68%
Other/unknown	24	30%	16	37%	8	22%
Subtotal	80	100%	43	100%	37	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	80%	2	67%	2	100%
Other/unknown	1	20%	1	33%	-	-
Subtotal	5	100%	3	100%	2	100%
Total	437		238		199	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 217. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Canton Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	253	79%	274	80%	298	80%	327	81%	352	81%
Females	67	21%	70	20%	75	20%	79	19%	85	19%
Age at diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	2	1%	2	1%	2	1%	3	1%	3	1%
20-24	2	1%	7	2%	11	3%	17	4%	26	6%
25-29	7	2%	14	4%	23	6%	25	6%	32	7%
30-34	16	5%	16	5%	17	5%	23	6%	26	6%
35-39	25	8%	26	8%	28	8%	28	7%	30	7%
40-44	63	20%	67	19%	73	20%	80	20%	82	19%
45-49	70	22%	73	21%	73	20%	77	19%	78	18%
50-54	54	17%	56	16%	60	16%	63	16%	66	15%
55-64	63	20%	65	19%	66	18%	70	17%	73	17%
65+	17	5%	17	5%	19	5%	19	5%	20	5%
Race/Ethnicity										
White, not Hispanic	229	72%	242	70%	261	70%	281	69%	301	69%
Black/African American, not Hispanic	76	24%	85	25%	92	25%	102	25%	111	25%
Hispanic/Latino	10	3%	11	3%	11	3%	12	3%	12	3%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
American Indian/Alaska Native	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
Unknown	3	1%	4	1%	7	2%	8	2%	9	2%
Race/Sex										
White, not Hispanic Males	187	58%	199	58%	216	58%	233	57%	250	57%
White, not Hispanic Females	42	13%	43	13%	45	12%	48	12%	51	12%
Black/African American, not Hispanic Males	53	17%	60	17%	65	17%	74	18%	81	19%
Black/African American, not Hispanic Females	23	7%	25	7%	27	7%	28	7%	30	7%
Hispanic/Latino Males	8	3%	9	3%	9	2%	10	2%	10	2%
Hispanic/Latina Females	2	1%	2	1%	2	1%	2	<1%	2	<1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	1	<1%	1	<1%	1	<1%	2	<1%	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	3	1%	4	1%	7	2%	8	2%	9	2%
Total	320		344		373		406		437	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 218. Reported persons living with a diagnosis of HIV infection by year and transmission category, Canton Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	175	69%	193	70%	209	70%	228	70%	247	70%
Injection drug use (IDU)	5	2%	6	2%	7	2%	7	2%	7	2%
Male-to-male sexual contact and IDU	8	3%	8	3%	9	3%	11	3%	12	3%
Heterosexual contact	13	5%	13	5%	16	5%	20	6%	21	6%
Other/unknown	52	21%	54	20%	57	19%	61	19%	65	18%
Subtotal	253	100%	274	100%	298	100%	327	100%	352	100%
Female adult or adolescent										
Injection drug use	6	10%	6	9%	6	9%	6	8%	8	10%
Heterosexual contact	36	58%	39	60%	43	61%	45	61%	48	60%
Other/unknown	20	32%	20	31%	21	30%	23	31%	24	30%
Subtotal	62	100%	65	100%	70	100%	74	100%	80	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	80%	4	80%	4	80%	4	80%	4	80%
Other/unknown	1	20%	1	20%	1	20%	1	20%	1	20%
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
Total	320		344		373		406		437	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 219. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Canton Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	4.6	15	88%	6.7	22	88%	7.9	26	84%	9.2	30	88%	7.9	26	81%
Females	*	2	12%	*	3	12%	1.5	5	16%	*	4	12%	1.8	6	19%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	3	12%	*	1	3%	*	4	12%	*	1	3%
20-24	*	2	12%	19.7	8	32%	27.0	11	35%	*	4	12%	22.1	9	28%
25-29	*	1	6%	*	1	4%	*	1	3%	*	3	9%	18.7	7	22%
30-34	*	3	18%	*	1	4%	*	3	10%	*	4	12%	*	2	6%
35-39	*	1	6%	*	3	12%	*	4	13%	13.4	5	15%	*	2	6%
40-44	14.2	6	35%	*	3	12%	*	2	6%	14.2	6	18%	*	2	6%
45-49	*	4	24%	*	3	12%	*	4	13%	*	1	3%	*	2	6%
50-54	*	-	-	*	2	8%	*	2	6%	*	4	12%	*	2	6%
55-64	*	-	-	*	1	4%	*	2	6%	*	3	9%	*	4	13%
65+	*	-	-	*	-	-	*	1	3%	*	-	-	*	1	3%
Race/Ethnicity															
White, not Hispanic	2.0	12	71%	2.1	13	52%	3.4	21	68%	3.4	21	62%	3.3	20	63%
Black/African American, not Hispanic	*	4	24%	31.6	10	40%	22.1	7	23%	31.6	10	29%	28.5	9	28%
Hispanic/Latino	*	1	6%	*	1	4%	*	-	-	*	1	3%	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-
Unknown	*	-	-	*	1	4%	*	3	10%	*	1	3%	*	2	6%
Race/Sex															
White, not Hispanic Males	3.7	11	65%	4.0	12	48%	6.4	19	61%	6.0	18	53%	5.7	17	53%
White, not Hispanic Females	*	1	6%	*	1	4%	*	2	6%	*	3	9%	*	3	9%
Black/African American, not Hispanic Males	*	3	18%	50.9	8	32%	31.8	5	16%	57.3	9	26%	44.5	7	22%
Black/African American, not Hispanic Females	*	1	6%	*	2	8%	*	2	6%	*	1	3%	*	2	6%
Hispanic/Latino Males	*	1	6%	*	1	4%	*	-	-	*	1	3%	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	4%	*	3	10%	*	1	3%	*	2	6%
Total	2.5	17		3.7	25		4.6	31		5.1	34		4.8	32	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 220. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Canton Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	10	67%	19	86%	17	65%	20	67%	19	73%
Injection drug use (IDU)	-	-	1	5%	2	8%	-	-	1	4%
Male-to-male sexual contact and IDU	-	-	-	-	1	4%	2	7%	1	4%
Heterosexual contact	1	7%	-	-	3	12%	4	13%	1	4%
Other/unknown	4	27%	2	9%	3	12%	4	13%	4	15%
Subtotal	15	100%	22	100%	26	100%	30	100%	26	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	2	33%
Heterosexual contact	2	100%	3	100%	4	80%	2	50%	3	50%
Other/unknown	-	-	-	-	1	20%	2	50%	1	17%
Subtotal	2	100%	3	100%	5	100%	4	100%	6	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	17		25		31		34		32	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 221. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Canton Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	7	88%	9	75%	11	73%	9	69%	13	76%	367	83%
Females	1	13%	3	25%	4	27%	4	31%	4	24%	74	17%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	6	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	1	<1%
20-24	-	-	1	8%	-	-	1	8%	1	6%	20	5%
25-29	-	-	-	-	-	-	2	15%	3	18%	52	12%
30-34	1	13%	1	8%	3	20%	1	8%	1	6%	91	21%
35-39	1	13%	2	17%	2	13%	2	15%	3	18%	93	21%
40-44	2	25%	3	25%	3	20%	3	23%	2	12%	74	17%
45-49	1	13%	3	25%	2	13%	2	15%	2	12%	41	9%
50-54	2	25%	2	17%	3	20%	-	-	2	12%	29	7%
55-64	1	13%	-	-	1	7%	2	15%	2	12%	24	5%
65+	-	-	-	-	1	7%	-	-	1	6%	10	2%
Race/Ethnicity												
White, not Hispanic	6	75%	9	75%	12	80%	5	38%	6	35%	326	74%
Black/African American, not Hispanic	2	25%	3	25%	2	13%	7	54%	9	53%	100	23%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	10	2%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	7%	1	8%	2	12%	5	1%
Race/Sex												
White, not Hispanic Males	5	63%	7	58%	8	53%	5	38%	5	29%	282	64%
White, not Hispanic Females	1	13%	2	17%	4	27%	-	-	1	6%	44	10%
Black/African American, not Hispanic Males	2	25%	2	17%	2	13%	3	23%	6	35%	72	16%
Black/African American, not Hispanic Females	-	-	1	8%	-	-	4	31%	3	18%	28	6%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	8	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	7%	1	8%	2	12%	5	1%
Total	8		12		15		13		17		441	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Canton Consortium 8

Table 222. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Canton Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	5	71%	8	89%	3	27%	8	89%	8	62%	258	71%
Injection drug use (IDU)	-	-	-	-	1	9%	-	-	1	8%	15	4%
Male-to-male sexual contact and IDU	-	-	-	-	1	9%	-	-	-	-	17	5%
Heterosexual contact	-	-	-	-	2	18%	-	-	3	23%	21	6%
Other/unknown	2	29%	1	11%	4	36%	1	11%	1	8%	54	15%
Subtotal	7	100%	9	100%	11	100%	9	100%	13	100%	365	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	25%	-	-	2	50%	14	20%
Heterosexual contact	1	100%	3	100%	3	75%	1	25%	2	50%	41	59%
Other/unknown	-	-	-	-	-	-	3	75%	-	-	15	21%
Subtotal	1	100%	3	100%	4	100%	4	100%	4	100%	70	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	6	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	8		12		15		13		17		441	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 223. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Lima Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	84.9	218	76%	111	77%	107	75%
Females	26.5	69	24%	34	23%	35	25%
Age at diagnosis (yr)							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	3	1%	-	-	3	2%
20-24	24.6	8	3%	7	5%	1	1%
25-29	47.4	14	5%	10	7%	4	3%
30-34	107.4	32	11%	19	13%	13	9%
35-39	91.6	27	9%	13	9%	14	10%
40-44	129.4	42	15%	22	15%	20	14%
45-49	153.2	56	20%	27	19%	29	20%
50-54	136.5	54	19%	26	18%	28	20%
55-64	57.1	39	14%	15	10%	24	17%
65+	15.7	12	4%	6	4%	6	4%
Race/Ethnicity							
White, not Hispanic	43.9	208	72%	106	73%	102	72%
Black/African American, not Hispanic	332.4	59	21%	29	20%	30	21%
Hispanic/Latino	100.9	13	5%	8	6%	5	4%
Asian/Pacific Islander	*	2	1%	1	1%	1	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	5	2%	1	1%	4	3%
Race/Sex							
White, not Hispanic Males	70.8	166	58%	86	59%	80	56%
White, not Hispanic Females	17.5	42	15%	20	14%	22	15%
Black/African American, not Hispanic Males	391.6	37	13%	17	12%	20	14%
Black/African American, not Hispanic Females	264.9	22	8%	12	8%	10	7%
Hispanic/Latino Males	133.2	9	3%	6	4%	3	2%
Hispanic/Latina Females	*	4	1%	2	1%	2	1%
Asian/Pacific Islander Males	*	2	1%	1	1%	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	5	2%	1	1%	4	3%
Total	55.5	287		145		142	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 224. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Lima Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	137	63%	67	60%	70	66%
Injection drug use (IDU)	7	3%	4	4%	3	3%
Male-to-male sexual contact and IDU	11	5%	8	7%	3	3%
Heterosexual contact	17	8%	5	5%	12	11%
Other/unknown	45	21%	27	24%	18	17%
Subtotal	217	100%	111	100%	106	100%
Female adult or adolescent						
Injection drug use	8	12%	3	9%	5	16%
Heterosexual contact	42	64%	20	59%	22	69%
Other/unknown	16	24%	11	32%	5	16%
Subtotal	66	100%	34	100%	32	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	100%	-	-	4	100%
Other/unknown	-	-	-	-	-	-
Subtotal	4	100%	-	-	4	100%
Total	287		145		142	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 225. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Lima Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	167	73%	177	74%	184	74%	203	75%	218	76%
Females	61	27%	63	26%	64	26%	67	25%	69	24%
Age at diagnosis (yr)										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	3	1%	3	1%	3	1%	3	1%	3	1%
20-24	1	<1%	1	<1%	1	<1%	4	1%	8	3%
25-29	6	3%	7	3%	10	4%	13	5%	14	5%
30-34	19	8%	23	10%	26	10%	29	11%	32	11%
35-39	20	9%	21	9%	22	9%	23	9%	27	9%
40-44	37	16%	38	16%	38	15%	40	15%	42	15%
45-49	50	22%	51	21%	51	21%	53	20%	56	20%
50-54	46	20%	49	20%	50	20%	54	20%	54	19%
55-64	34	15%	35	15%	35	14%	39	14%	39	14%
65+	12	5%	12	5%	12	5%	12	4%	12	4%
Race/Ethnicity										
White, not Hispanic	161	71%	171	71%	177	71%	196	73%	208	72%
Black/African American, not Hispanic	51	22%	52	22%	54	22%	56	21%	59	21%
Hispanic/Latino	11	5%	12	5%	12	5%	12	4%	13	5%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	2	1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	4	2%	4	2%	4	2%	5	2%	5	2%
Race/Sex										
White, not Hispanic Males	126	55%	134	56%	139	56%	155	57%	166	58%
White, not Hispanic Females	35	15%	37	15%	38	15%	41	15%	42	15%
Black/African American, not Hispanic Males	30	13%	31	13%	33	13%	35	13%	37	13%
Black/African American, not Hispanic Females	21	9%	21	9%	21	8%	21	8%	22	8%
Hispanic/Latino Males	7	3%	8	3%	8	3%	8	3%	9	3%
Hispanic/Latina Females	4	2%	4	2%	4	2%	4	1%	4	1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	2	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	4	2%	4	2%	4	2%	5	2%	5	2%
Total	228		240		248		270		287	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 226. Reported persons living with a diagnosis of HIV infection by year and transmission category, Lima Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	112	67%	116	66%	122	67%	129	64%	137	63%
Injection drug use (IDU)	7	4%	7	4%	7	4%	7	3%	7	3%
Male-to-male sexual contact and IDU	10	6%	10	6%	10	5%	11	5%	11	5%
Heterosexual contact	12	7%	16	9%	16	9%	17	8%	17	8%
Other/unknown	25	15%	27	15%	28	15%	38	19%	45	21%
Subtotal	166	100%	176	100%	183	100%	202	100%	217	100%
Female adult or adolescent										
Injection drug use	6	10%	7	12%	8	13%	8	13%	8	12%
Heterosexual contact	37	64%	38	63%	38	62%	41	64%	42	64%
Other/unknown	15	26%	15	25%	15	25%	15	23%	16	24%
Subtotal	58	100%	60	100%	61	100%	64	100%	66	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	100%	4	100%	4	100%	4	100%	4	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	4	100%	4	100%	4	100%	4	100%	4	100%
Total	228		240		248		270		287	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 227. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Lima Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	5.5	14	93%	4.7	12	75%	3.5	9	90%	7.4	19	86%	5.8	15	88%
Females	*	1	7%	*	4	25%	*	1	10%	*	3	14%	*	2	12%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	3	20%	*	1	6%	*	1	10%	*	4	18%	15.4	5	29%
25-29	*	4	27%	*	3	19%	16.9	5	50%	*	4	18%	*	1	6%
30-34	*	2	13%	*	3	19%	*	-	-	*	2	9%	*	3	18%
35-39	*	2	13%	*	1	6%	*	1	10%	*	1	5%	*	3	18%
40-44	*	1	7%	*	1	6%	*	1	10%	*	2	9%	15.4	5	29%
45-49	*	1	7%	13.7	5	31%	*	1	10%	*	4	18%	*	-	-
50-54	*	1	7%	*	1	6%	*	-	-	*	3	14%	*	-	-
55-64	*	1	7%	*	1	6%	*	1	10%	*	2	9%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity															
White, not Hispanic	2.3	11	73%	2.3	11	69%	1.5	7	70%	4.0	19	86%	2.5	12	71%
Black/African American, not Hispanic	*	2	13%	*	3	19%	*	3	30%	*	2	9%	*	3	18%
Hispanic/Latino	*	1	7%	*	2	13%	*	-	-	*	-	-	*	1	6%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	6%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	7%	*	-	-	*	-	-	*	1	5%	*	-	-
Race/Sex															
White, not Hispanic Males	4.3	10	67%	3.8	9	56%	2.6	6	60%	6.8	16	73%	4.7	11	65%
White, not Hispanic Females	*	1	7%	*	2	13%	*	1	10%	*	3	14%	*	1	6%
Black/African American, not Hispanic Males	*	2	13%	*	2	13%	*	3	30%	*	2	9%	*	2	12%
Black/African American, not Hispanic Females	*	-	-	*	1	6%	*	-	-	*	-	-	*	1	6%
Hispanic/Latino Males	*	1	7%	*	1	6%	*	-	-	*	-	-	*	1	6%
Hispanic/Latina Females	*	-	-	*	1	6%	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	1	6%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	1	7%	*	-	-	*	-	-	*	1	5%	*	-	-
Total	2.9	15		3.1	16		1.9	10		4.3	22		3.3	17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 228. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Lima Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	13	93%	4	33%	8	89%	7	37%	8	53%
Injection drug use (IDU)	-	-	2	17%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	5%	-	-
Heterosexual contact	1	7%	4	33%	-	-	1	5%	-	-
Other/unknown	-	-	2	17%	1	11%	10	53%	7	47%
Subtotal	14	100%	12	100%	9	100%	19	100%	15	100%
Female adult or adolescent										
Injection drug use	-	-	2	50%	1	100%	-	-	-	-
Heterosexual contact	1	100%	1	25%	-	-	3	100%	1	50%
Other/unknown	-	-	1	25%	-	-	-	-	1	50%
Subtotal	1	100%	4	100%	1	100%	3	100%	2	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	15		16		10		22		17	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 229. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Lima Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	7	70%	7	64%	5	42%	11	85%	6	75%	263	82%
Females	3	30%	4	36%	7	58%	2	15%	2	25%	57	18%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	5	2%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	3	1%
20-24	-	-	-	-	1	8%	-	-	-	-	7	2%
25-29	-	-	2	18%	1	8%	4	31%	3	38%	50	16%
30-34	2	20%	3	27%	1	8%	2	15%	1	13%	58	18%
35-39	-	-	1	9%	-	-	-	-	1	13%	52	16%
40-44	-	-	1	9%	3	25%	1	8%	-	-	45	14%
45-49	2	20%	2	18%	4	33%	4	31%	1	13%	45	14%
50-54	5	50%	1	9%	1	8%	2	15%	1	13%	30	9%
55-64	1	10%	1	9%	1	8%	-	-	1	13%	20	6%
65+	-	-	-	-	-	-	-	-	-	-	5	2%
Race/Ethnicity												
White, not Hispanic	7	70%	7	64%	7	58%	11	85%	6	75%	260	81%
Black/African American, not Hispanic	2	20%	4	36%	4	33%	-	-	2	25%	45	14%
Hispanic/Latino	-	-	-	-	-	-	-	-	-	-	9	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	10%	-	-	1	8%	2	15%	-	-	5	2%
Race/Sex												
White, not Hispanic Males	6	60%	5	45%	3	25%	9	69%	5	63%	225	70%
White, not Hispanic Females	1	10%	2	18%	4	33%	2	15%	1	13%	35	11%
Black/African American, not Hispanic Males	1	10%	2	18%	2	17%	-	-	1	13%	28	9%
Black/African American, not Hispanic Females	1	10%	2	18%	2	17%	-	-	1	13%	17	5%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	-	-	6	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	3	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	1	10%	-	-	1	8%	2	15%	-	-	5	2%
Total	10		11		12		13		8		320	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Lima Consortium 9A

Table 230. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Lima Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	5	71%	2	29%	4	80%	6	55%	4	67%	176	67%
Injection drug use (IDU)	-	-	1	14%	1	20%	-	-	-	-	10	4%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	10	4%
Heterosexual contact	1	14%	2	29%	-	-	1	9%	-	-	19	7%
Other/unknown	1	14%	2	29%	-	-	4	36%	2	33%	48	18%
Subtotal	7	100%	7	100%	5	100%	11	100%	6	100%	263	100%
Female adult or adolescent												
Injection drug use	1	33%	1	25%	1	14%	-	-	-	-	8	15%
Heterosexual contact	2	67%	2	50%	4	57%	2	100%	1	50%	37	71%
Other/unknown	-	-	1	25%	2	29%	-	-	1	50%	7	13%
Subtotal	3	100%	4	100%	7	100%	2	100%	2	100%	52	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	80%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	20%
Subtotal	-	-	-	-	-	-	-	-	-	-	5	100%
Total	10		11		12		13		8		320	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 231. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Mansfield Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	78.2	233	77%	97	72%	136	80%
Females	23.7	71	23%	37	28%	34	20%
Age at diagnosis (yr)							
<13	*	1	<1%	1	1%	-	-
13-14	*	1	<1%	1	1%	-	-
15-19	*	2	1%	1	1%	1	1%
20-24	40.9	15	5%	11	8%	4	2%
25-29	65.7	22	7%	16	12%	6	4%
30-34	71.6	25	8%	12	9%	13	8%
35-39	86.8	30	10%	15	11%	15	9%
40-44	86.2	33	11%	17	13%	16	9%
45-49	137.5	58	19%	20	15%	38	22%
50-54	107.7	50	16%	13	10%	37	22%
55-64	60.7	50	16%	22	16%	28	16%
65+	18.0	17	6%	5	4%	12	7%
Race/Ethnicity							
White, not Hispanic	41.0	223	73%	101	75%	122	72%
Black/African American, not Hispanic	227.3	58	19%	23	17%	35	21%
Hispanic/Latino	101.6	15	5%	7	5%	8	5%
Asian/Pacific Islander	*	1	<1%	1	1%	-	-
American Indian/Alaska Native	*	1	<1%	-	-	1	1%
Unknown	*	6	2%	2	1%	4	2%
Race/Sex							
White, not Hispanic Males	66.3	178	59%	77	57%	101	59%
White, not Hispanic Females	16.3	45	15%	24	18%	21	12%
Black/African American, not Hispanic Males	251.5	39	13%	12	9%	27	16%
Black/African American, not Hispanic Females	189.8	19	6%	11	8%	8	5%
Hispanic/Latino Males	166.9	13	4%	7	5%	6	4%
Hispanic/Latina Females	*	2	1%	-	-	2	1%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	1	<1%	1	1%	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	1%
Unknown	*	6	2%	2	1%	4	2%
Total	50.9	304		134		170	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 232. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Mansfield Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	152	66%	56	60%	96	71%
Injection drug use (IDU)	6	3%	1	1%	5	4%
Male-to-male sexual contact and IDU	14	6%	3	3%	11	8%
Heterosexual contact	16	7%	7	7%	9	7%
Other/unknown	41	18%	27	29%	14	10%
Subtotal	229	100%	94	100%	135	100%
Female adult or adolescent						
Injection drug use	4	6%	1	3%	3	9%
Heterosexual contact	49	71%	26	72%	23	70%
Other/unknown	16	23%	9	25%	7	21%
Subtotal	69	100%	36	100%	33	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	67%	2	50%	2	100%
Other/unknown	2	33%	2	50%	-	-
Subtotal	6	100%	4	100%	2	100%
Total	304		134		170	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 233. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Mansfield Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	190	77%	199	76%	211	77%	219	77%	233	77%
Females	57	23%	62	24%	64	23%	66	23%	71	23%
Age at diagnosis (yr)										
<13	-	-	-	-	1	<1%	1	<1%	1	<1%
13-14	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
15-19	1	<1%	1	<1%	1	<1%	2	1%	2	1%
20-24	2	1%	4	2%	8	3%	10	4%	15	5%
25-29	13	5%	14	5%	15	5%	19	7%	22	7%
30-34	21	9%	23	9%	23	8%	24	8%	25	8%
35-39	21	9%	24	9%	26	9%	28	10%	30	10%
40-44	30	12%	31	12%	32	12%	32	11%	33	11%
45-49	51	21%	52	20%	55	20%	55	19%	58	19%
50-54	46	19%	48	18%	50	18%	50	18%	50	16%
55-64	46	19%	47	18%	47	17%	47	16%	50	16%
65+	15	6%	16	6%	16	6%	16	6%	17	6%
Race/Ethnicity										
White, not Hispanic	182	74%	190	73%	201	73%	208	73%	223	73%
Black/African American, not Hispanic	49	20%	53	20%	55	20%	58	20%	58	19%
Hispanic/Latino	11	4%	11	4%	12	4%	12	4%	15	5%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	5	2%	6	2%	6	2%	6	2%	6	2%
Race/Sex										
White, not Hispanic Males	145	59%	152	58%	161	59%	167	59%	178	59%
White, not Hispanic Females	37	15%	38	15%	40	15%	41	14%	45	15%
Black/African American, not Hispanic Males	33	13%	35	13%	37	13%	39	14%	39	13%
Black/African American, not Hispanic Females	16	6%	18	7%	18	7%	19	7%	19	6%
Hispanic/Latino Males	9	4%	9	3%	10	4%	10	4%	13	4%
Hispanic/Latina Females	2	1%	2	1%	2	1%	2	1%	2	1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	<1%	1	<1%	1	<1%	1	<1%
Unknown	5	2%	6	2%	6	2%	6	2%	6	2%
Total	247		261		275		285		304	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 234. Reported persons living with a diagnosis of HIV infection by year and transmission category, Mansfield Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	122	66%	127	65%	135	65%	142	66%	152	66%
Injection drug use (IDU)	6	3%	6	3%	6	3%	6	3%	6	3%
Male-to-male sexual contact and IDU	14	8%	14	7%	14	7%	14	7%	14	6%
Heterosexual contact	11	6%	12	6%	13	6%	13	6%	16	7%
Other/unknown	33	18%	36	18%	39	19%	40	19%	41	18%
Subtotal	186	100%	195	100%	207	100%	215	100%	229	100%
Female adult or adolescent										
Injection drug use	3	5%	3	5%	3	5%	3	5%	4	6%
Heterosexual contact	39	70%	44	72%	45	73%	46	72%	49	71%
Other/unknown	14	25%	14	23%	14	23%	15	23%	16	23%
Subtotal	56	100%	61	100%	62	100%	64	100%	69	100%
Child (<13 yrs at diagnosis)										
Perinatal	4	80%	4	80%	4	67%	4	67%	4	67%
Other/unknown	1	20%	1	20%	2	33%	2	33%	2	33%
Subtotal	5	100%	5	100%	6	100%	6	100%	6	100%
Total	247		261		275		285		304	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 235. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Mansfield Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	4.0	12	71%	3.0	9	75%	4.0	12	86%	2.7	8	80%	4.7	14	74%
Females	1.7	5	29%	*	3	25%	*	2	14%	*	2	20%	1.7	5	26%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	1	7%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	1	8%	*	2	14%	*	2	20%	*	-	-
20-24	*	2	12%	*	1	8%	*	2	14%	*	3	30%	16.4	6	32%
25-29	*	2	12%	*	-	-	*	1	7%	*	2	20%	*	3	16%
30-34	*	3	18%	*	2	17%	*	2	14%	*	2	20%	*	1	5%
35-39	*	3	18%	*	2	17%	*	1	7%	*	1	10%	*	1	5%
40-44	13.1	5	29%	*	2	17%	*	3	21%	*	-	-	*	1	5%
45-49	*	1	6%	*	2	17%	*	1	7%	*	-	-	*	3	16%
50-54	*	-	-	*	1	8%	*	1	7%	*	-	-	*	-	-
55-64	*	1	6%	*	1	8%	*	-	-	*	-	-	*	3	16%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	5%
Race/Ethnicity															
White, not Hispanic	2.2	12	71%	1.5	8	67%	2.0	11	79%	1.3	7	70%	2.8	15	79%
Black/African American, not Hispanic	19.6	5	29%	*	3	25%	*	2	14%	*	3	30%	*	-	-
Hispanic/Latino	*	-	-	*	-	-	*	1	7%	*	-	-	*	3	16%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	1	5%
American Indian/Alaska Native	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Sex															
White, not Hispanic Males	2.6	7	41%	2.6	7	58%	3.4	9	64%	2.2	6	60%	4.1	11	58%
White, not Hispanic Females	1.8	5	29%	*	1	8%	*	2	14%	*	1	10%	*	4	21%
Black/African American, not Hispanic Males	32.2	5	29%	*	2	17%	*	2	14%	*	2	20%	*	-	-
Black/African American, not Hispanic Females	*	-	-	*	1	8%	*	-	-	*	1	10%	*	-	-
Hispanic/Latino Males	*	-	-	*	-	-	*	1	7%	*	-	-	*	3	16%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	5%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	2.8	17		2.0	12		2.3	14		1.7	10		3.2	19	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 236. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Mansfield Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	9	75%	4	44%	8	67%	7	88%	10	71%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	1	8%	1	11%	1	8%	-	-	3	21%
Other/unknown	2	17%	4	44%	3	25%	1	13%	1	7%
Subtotal	12	100%	9	100%	12	100%	8	100%	14	100%
Female adult or adolescent										
Injection drug use	-	-	-	-	-	-	-	-	1	20%
Heterosexual contact	2	40%	3	100%	1	100%	1	50%	3	60%
Other/unknown	3	60%	-	-	-	-	1	50%	1	20%
Subtotal	5	100%	3	100%	1	100%	2	100%	5	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-
Total	17		12		14		10		19	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 237. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Mansfield Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	6	67%	11	79%	7	88%	3	60%	6	67%	354	85%
Females	3	33%	3	21%	1	13%	2	40%	3	33%	61	15%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	7	2%
13-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	-	-	1	11%	2	<1%
20-24	2	22%	1	7%	-	-	1	20%	1	11%	21	5%
25-29	1	11%	1	7%	1	13%	-	-	1	11%	60	14%
30-34	-	-	1	7%	-	-	1	20%	1	11%	65	16%
35-39	3	33%	-	-	2	25%	3	60%	1	11%	95	23%
40-44	1	11%	5	36%	3	38%	-	-	2	22%	71	17%
45-49	2	22%	5	36%	-	-	-	-	1	11%	52	13%
50-54	-	-	-	-	2	25%	-	-	-	-	17	4%
55-64	-	-	1	7%	-	-	-	-	1	11%	19	5%
65+	-	-	-	-	-	-	-	-	-	-	5	1%
Race/Ethnicity												
White, not Hispanic	7	78%	9	64%	5	63%	4	80%	8	89%	345	83%
Black/African American, not Hispanic	2	22%	4	29%	2	25%	1	20%	1	11%	55	13%
Hispanic/Latino	-	-	-	-	1	13%	-	-	-	-	11	3%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native	-	-	1	7%	-	-	-	-	-	-	2	<1%
Unknown	-	-	-	-	-	-	-	-	-	-	2	<1%
Race/Sex												
White, not Hispanic Males	4	44%	8	57%	4	50%	3	60%	6	67%	301	73%
White, not Hispanic Females	3	33%	1	7%	1	13%	1	20%	2	22%	44	11%
Black/African American, not Hispanic Males	2	22%	3	21%	2	25%	-	-	-	-	42	10%
Black/African American, not Hispanic Females	-	-	1	7%	-	-	1	20%	1	11%	13	3%
Hispanic/Latino Males	-	-	-	-	1	13%	-	-	-	-	9	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	1	7%	-	-	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-	-	-	-	-	2	<1%
Total	9		14		8		5		9		415	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Mansfield Consortium 9B

Table 238. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Mansfield Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	5	83%	9	82%	4	57%	3	100%	4	67%	239	68%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	20	6%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	30	9%
Heterosexual contact	-	-	-	-	1	14%	-	-	-	-	15	4%
Other/unknown	1	17%	2	18%	2	29%	-	-	2	33%	46	13%
Subtotal	6	100%	11	100%	7	100%	3	100%	6	100%	350	100%
Female adult or adolescent												
Injection drug use	1	33%	-	-	-	-	-	-	-	-	8	14%
Heterosexual contact	1	33%	3	100%	1	100%	-	-	2	67%	39	67%
Other/unknown	1	33%	-	-	-	-	2	100%	1	33%	11	19%
Subtotal	3	100%	3	100%	1	100%	2	100%	3	100%	58	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	71%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	29%
Subtotal	-	-	-	-	-	-	-	-	-	-	7	100%
Total	9		14		8		5		9		415	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 239. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and selected characteristics, Southeast Part B

Characteristic	Living with a diagnosis of HIV infection - 2011			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	79.2	335	76%	151	72%	184	81%
Females	24.4	104	24%	60	28%	44	19%
Age at diagnosis (yr)							
<13	*	2	<1%	1	<1%	1	<1%
13-14	*	-	-	-	-	-	-
15-19	*	2	<1%	-	-	2	1%
20-24	17.8	11	3%	10	5%	1	<1%
25-29	34.9	17	4%	11	5%	6	3%
30-34	62.1	31	7%	20	9%	11	5%
35-39	92.8	46	10%	29	14%	17	7%
40-44	122.9	67	15%	33	16%	34	15%
45-49	159.6	94	21%	31	15%	63	28%
50-54	114.2	74	17%	27	13%	47	21%
55-64	66.6	77	18%	40	19%	37	16%
65+	13.7	18	4%	9	4%	9	4%
Race/Ethnicity							
White, not Hispanic	46.8	374	85%	175	83%	199	87%
Black/African American, not Hispanic	200.7	44	10%	23	11%	21	9%
Hispanic/Latino	112.3	9	2%	5	2%	4	2%
Asian/Pacific Islander	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	11	3%	8	4%	3	1%
Race/Sex							
White, not Hispanic Males	73.5	290	66%	127	60%	163	71%
White, not Hispanic Females	20.8	84	19%	48	23%	36	16%
Black/African American, not Hispanic Males	211.1	30	7%	15	7%	15	7%
Black/African American, not Hispanic Females	181.5	14	3%	8	4%	6	3%
Hispanic/Latino Males	143.0	6	1%	4	2%	2	1%
Hispanic/Latina Females	*	3	1%	1	<1%	2	1%
Asian/Pacific Islander Males	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Unknown	*	11	3%	8	4%	3	1%
Total	51.7	439		211		228	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 240. Reported persons living with a diagnosis of HIV infection as of December 31, 2011, by current disease status and transmission category, Southeast Part B

Transmission Category	Living with a diagnosis of HIV infection - 2011		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	217	66%	92	61%	125	70%
Injection drug use (IDU)	13	4%	4	3%	9	5%
Male-to-male sexual contact and IDU	11	3%	4	3%	7	4%
Heterosexual contact	29	9%	10	7%	19	11%
Other/unknown	60	18%	41	27%	19	11%
Subtotal	330	100%	151	100%	179	100%
Female adult or adolescent						
Injection drug use	9	9%	6	11%	3	7%
Heterosexual contact	59	59%	27	47%	32	74%
Other/unknown	32	32%	24	42%	8	19%
Subtotal	100	100%	57	100%	43	100%
Child (<13 yrs at diagnosis)						
Perinatal	6	67%	3	100%	3	50%
Other/unknown	3	33%	-	-	3	50%
Subtotal	9	100%	3	100%	6	100%
Total	439		211		228	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection by current disease status categorizes cases by those diagnosed with HIV (not AIDS) and those diagnosed with AIDS as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 241. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Southeast Part B, 2007-2011

Characteristic	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	261	75%	277	76%	298	76%	317	77%	335	76%
Females	85	25%	88	24%	92	24%	96	23%	104	24%
Age at diagnosis (yr)										
<13	1	<1%	1	<1%	2	1%	2	<1%	2	<1%
13-14	-	-	-	-	-	-	-	-	-	-
15-19	2	1%	2	1%	2	1%	2	<1%	2	<1%
20-24	3	1%	4	1%	5	1%	6	1%	11	3%
25-29	6	2%	8	2%	11	3%	13	3%	17	4%
30-34	18	5%	20	5%	23	6%	29	7%	31	7%
35-39	25	7%	34	9%	39	10%	42	10%	46	10%
40-44	53	15%	55	15%	59	15%	66	16%	67	15%
45-49	87	25%	87	24%	90	23%	92	22%	94	21%
50-54	66	19%	66	18%	67	17%	69	17%	74	17%
55-64	69	20%	71	19%	75	19%	75	18%	77	18%
65+	16	5%	17	5%	17	4%	17	4%	18	4%
Race/Ethnicity										
White, not Hispanic	296	86%	311	85%	330	85%	349	85%	374	85%
Black/African American, not Hispanic	35	10%	38	10%	41	11%	43	10%	44	10%
Hispanic/Latino	5	1%	5	1%	8	2%	9	2%	9	2%
Asian/Pacific Islander	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-
Unknown	9	3%	10	3%	10	3%	11	3%	11	3%
Race/Sex										
White, not Hispanic Males	227	66%	240	66%	256	66%	273	66%	290	66%
White, not Hispanic Females	69	20%	71	19%	74	19%	76	18%	84	19%
Black/African American, not Hispanic Males	24	7%	26	7%	29	7%	29	7%	30	7%
Black/African American, not Hispanic Females	11	3%	12	3%	12	3%	14	3%	14	3%
Hispanic/Latino Males	3	1%	3	1%	5	1%	6	1%	6	1%
Hispanic/Latina Females	2	1%	2	1%	3	1%	3	1%	3	1%
Asian/Pacific Islander Males	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-
Unknown	9	3%	10	3%	10	3%	11	3%	11	3%
Total	346		365		390		413		439	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 242. Reported persons living with a diagnosis of HIV infection by year and transmission category, Southeast Part B, 2007-2011

Transmission Category	Living with a Diagnosis of HIV Infection									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	162	63%	176	64%	191	65%	206	66%	217	66%
Injection drug use (IDU)	13	5%	13	5%	13	4%	13	4%	13	4%
Male-to-male sexual contact and IDU	9	4%	9	3%	9	3%	11	4%	11	3%
Heterosexual contact	27	11%	27	10%	28	10%	28	9%	29	9%
Other/unknown	46	18%	48	18%	52	18%	54	17%	60	18%
Subtotal	257	100%	273	100%	293	100%	312	100%	330	100%
Female adult or adolescent										
Injection drug use	6	7%	7	8%	7	8%	7	8%	9	9%
Heterosexual contact	49	60%	51	61%	54	61%	55	60%	59	59%
Other/unknown	26	32%	26	31%	27	31%	30	33%	32	32%
Subtotal	81	100%	84	100%	88	100%	92	100%	100	100%
Child (<13 yrs at diagnosis)										
Perinatal	5	63%	5	63%	6	67%	6	67%	6	67%
Other/unknown	3	38%	3	38%	3	33%	3	33%	3	33%
Subtotal	8	100%	8	100%	9	100%	9	100%	9	100%
Total	346		365		390		413		439	

Notes: Living with a diagnosis of HIV infection by year (2007-2011) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2011. Living with a diagnosis of HIV infection includes those diagnosed with HIV (not AIDS) and those diagnosed with AIDS by current disease status as of December 31, 2011.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 243. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Southeast Part B, 2007-2011

Characteristic	Diagnosis of HIV Infection by Year														
	2007			2008			2009			2010			2011		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	2.4	10	77%	4.3	18	82%	5.7	24	86%	4.3	18	82%	4.5	19	70%
Females	*	3	23%	*	4	18%	*	4	14%	*	4	18%	1.9	8	30%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	1	4%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	1	5%	*	-	-	*	-	-	*	1	4%
20-24	*	1	8%	*	2	9%	*	3	11%	*	2	9%	13.0	8	30%
25-29	*	2	15%	*	2	9%	*	4	14%	*	3	14%	*	1	4%
30-34	*	1	8%	16.0	8	36%	*	4	14%	10.0	5	23%	*	1	4%
35-39	*	1	8%	*	4	18%	12.1	6	21%	10.1	5	23%	10.1	5	19%
40-44	*	2	15%	*	2	9%	*	3	11%	9.2	5	23%	*	1	4%
45-49	8.5	5	38%	*	-	-	*	2	7%	*	1	5%	8.5	5	19%
50-54	*	-	-	*	2	9%	*	2	7%	*	1	5%	*	2	7%
55-64	*	1	8%	*	-	-	*	3	11%	*	-	-	*	2	7%
65+	*	-	-	*	1	5%	*	-	-	*	-	-	*	1	4%
Race/Ethnicity															
White, not Hispanic	1.6	13	100%	2.3	18	82%	2.8	22	79%	2.3	18	82%	3.3	26	96%
Black/African American, not Hispanic	*	-	-	*	3	14%	*	3	11%	*	2	9%	*	1	4%
Hispanic/Latino	*	-	-	*	-	-	*	3	11%	*	1	5%	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	5%	*	-	-	*	1	5%	*	-	-
Race/Sex															
White, not Hispanic Males	2.5	10	77%	3.8	15	68%	4.8	19	68%	4.1	16	73%	4.6	18	67%
White, not Hispanic Females	*	3	23%	*	3	14%	*	3	11%	*	2	9%	2.0	8	30%
Black/African American, not Hispanic Males	*	-	-	*	2	9%	*	3	11%	*	-	-	*	1	4%
Black/African American, not Hispanic Females	*	-	-	*	1	5%	*	-	-	*	2	9%	*	-	-
Hispanic/Latino Males	*	-	-	*	-	-	*	2	7%	*	1	5%	*	-	-
Hispanic/Latina Females	*	-	-	*	-	-	*	1	4%	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	1	5%	*	-	-	*	1	5%	*	-	-
Total	1.5	13		2.6	22		3.3	28		2.6	22		3.2	27	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2011 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 244. Reported diagnoses of HIV infection by year of diagnosis and transmission category, Southeast Part B, 2007-2011

Transmission Category	Diagnosis of HIV Infection by Year									
	2007		2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	6	60%	16	89%	18	78%	15	83%	11	58%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	6%	-	-
Heterosexual contact	1	10%	-	-	1	4%	-	-	1	5%
Other/unknown	3	30%	2	11%	4	17%	2	11%	7	37%
Subtotal	10	100%	18	100%	23	100%	18	100%	19	100%
Female adult or adolescent										
Injection drug use	-	-	1	25%	-	-	-	-	2	25%
Heterosexual contact	2	67%	2	50%	3	75%	1	25%	4	50%
Other/unknown	1	33%	1	25%	1	25%	3	75%	2	25%
Subtotal	3	100%	4	100%	4	100%	4	100%	8	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-
Total	13		22		28		22		27	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2007-2010) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 245. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Southeast Part B, 2007-2011

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex												
Males	14	88%	6	60%	10	67%	8	73%	9	82%	446	86%
Females	2	13%	4	40%	5	33%	3	27%	2	18%	71	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	1	7%	-	-	-	-	5	1%
13-14	-	-	-	-	-	-	-	-	-	-	3	1%
15-19	-	-	-	-	-	-	-	-	-	-	5	1%
20-24	1	6%	-	-	1	7%	1	9%	1	9%	12	2%
25-29	-	-	1	10%	2	13%	2	18%	1	9%	63	12%
30-34	1	6%	2	20%	-	-	2	18%	1	9%	101	20%
35-39	1	6%	3	30%	3	20%	2	18%	2	18%	100	19%
40-44	7	44%	2	20%	4	27%	2	18%	1	9%	116	22%
45-49	3	19%	2	20%	2	13%	1	9%	2	18%	55	11%
50-54	2	13%	-	-	1	7%	1	9%	1	9%	27	5%
55-64	1	6%	-	-	1	7%	-	-	2	18%	26	5%
65+	-	-	-	-	-	-	-	-	-	-	4	1%
Race/Ethnicity												
White, not Hispanic	14	88%	10	100%	12	80%	10	91%	11	100%	473	91%
Black/African American, not Hispanic	-	-	-	-	1	7%	1	9%	-	-	32	6%
Hispanic/Latino	1	6%	-	-	1	7%	-	-	-	-	5	1%
Asian/Pacific Islander	1	6%	-	-	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	7%	-	-	-	-	5	1%
Race/Sex												
White, not Hispanic Males	12	75%	6	60%	8	53%	8	73%	9	82%	412	80%
White, not Hispanic Females	2	13%	4	40%	4	27%	2	18%	2	18%	61	12%
Black/African American, not Hispanic Males	-	-	-	-	1	7%	-	-	-	-	23	4%
Black/African American, not Hispanic Females	-	-	-	-	-	-	1	9%	-	-	9	2%
Hispanic/Latino Males	1	6%	-	-	-	-	-	-	-	-	4	1%
Hispanic/Latina Females	-	-	-	-	1	7%	-	-	-	-	1	<1%
Asian/Pacific Islander Males	1	6%	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	1	7%	-	-	-	-	5	1%
Total	16		10		15		11		11		517	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

Ryan White Part B Southeast Consortium 9C

Table 246. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission category, Southeast Part B, 2007-2011

Transmission Category	Year of AIDS Diagnosis										Cumulative AIDS	
	2007		2008		2009		2010		2011		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	7	50%	4	67%	8	89%	6	75%	4	44%	297	68%
Injection drug use (IDU)	1	7%	1	17%	-	-	-	-	-	-	16	4%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	21	5%
Heterosexual contact	2	14%	1	17%	1	11%	-	-	-	-	22	5%
Other/unknown	4	29%	-	-	-	-	2	25%	5	56%	81	19%
Subtotal	14	100%	6	100%	9	100%	8	100%	9	100%	437	100%
Female adult or adolescent												
Injection drug use	1	50%	-	-	-	-	-	-	-	-	11	16%
Heterosexual contact	1	50%	4	100%	4	80%	1	33%	2	100%	41	59%
Other/unknown	-	-	-	-	1	20%	2	67%	-	-	18	26%
Subtotal	2	100%	4	100%	5	100%	3	100%	2	100%	70	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	1	100%	-	-	-	-	4	40%
Other/unknown	-	-	-	-	-	-	-	-	-	-	6	60%
Subtotal	-	-	-	-	1	100%	-	-	-	-	10	100%
Total	16		10		15		11		11		517	

Note: AIDS diagnoses by year (2007-2011) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2011.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through September 30, 2012.

This page is intentionally blank.